

Quinn, Amie L.

2007

The impacts of agricultural chemicals and temperature on the physiological stress response in fish

Department of Biological Studies

<https://hdl.handle.net/10133/676>

Downloaded from OPUS, University of Lethbridge Research Repository

THE IMPACTS OF AGRICULTURAL CHEMICALS AND TEMPERATURE ON THE
PHYSIOLOGICAL STRESS RESPONSE IN FISH

AMIE L. QUINN
Bachelor of Science, University of Lethbridge, 2005

A Thesis
Submitted to the School of Graduate Studies
of the University of Lethbridge
in Partial Fulfilment of the
Requirements for the Degree

MASTER OF SCIENCE

Department of Biological Sciences
University of Lethbridge
LETHBRIDGE, ALBERTA, CANADA

© Amie L. Quinn 2007

Abstract

Fish are exposed to multiple stressors in their environment. The interactive effects of pesticide exposure and increased temperature on the physiological stress response were investigated in a comparative field study with cold-water (whitefish, *Prosopium williamsoni*) and cool-water (sucker, *Catostomus*) fish from the Oldman River, Alberta, Canada, and in a laboratory study with rainbow trout, *Oncorhynchus mykiss*. Physiological stress indicators were measured, and exposure to pesticides was estimated using acetylcholinesterase (AChE) inhibition. Species-specific differences in AChE activities and responses of the physiological stress axis were detected in whitefish and suckers, suggesting that whitefish are a more sensitive species to temperature and pesticide stress. *In vivo* Dimethoate exposure inhibited AChE activity in various tissues and disrupted the physiological stress response. Commercial Dimethoate, *in vitro*, caused a decrease in viability and cortisol secretion while pure grade Dimethoate did not. The results from this study can be used in predictions of fish vulnerability to stress.

Acknowledgements

First of all I would like to thank my supervisor, Dr. Alice Hontela, for her never ending support, encouragement and time throughout this project. Her willingness to bring me into her project and provide guidance and advice were key to the successful completion of this thesis. I would also like to extend my deep thanks to the other members of my committee: Dr. Rasmussen, Dr. Byrne, and Dr. Rood for their comments and suggestions to further develop my thesis.

This paper would not have been possible without many others as well including, Caitlin Friesen, Ian Harper, Lindsay Akins, Lana Miller, Paul Dawson and Rheana Flitton for their help in the field and in the lab. Further thanks are also due to Wendell Koning from Alberta Environment for data on pesticide concentrations and temperature in the Oldman River, David Humphries for conducting dimethoate analysis on my water samples, Bruce McMullin for supplies, Tom Kevder for the donation of Cygon (dimethoate), and the Allison Creek Trout Hatchery for supplying my fish. A key thanks for the project also goes out to Alberta Ingenuity Centre for Water Research and the Canadian Water Network who funded the activity as well as to the University of Lethbridge for a wide range of services.

Lastly, I would like to thank my husband, Kyle Quinn, who supported me from the moment I said I wanted to do my Masters despite the personal challenges it entailed.

Table of Contents

| | |
|---|------|
| Title Page | i |
| Approval/Signature Page | ii |
| Abstract | iii |
| Acknowledgements | iv |
| Table of Contents | v |
| List of Tables | vii |
| List of Figures | viii |
| List of Abbreviations | ix |
| Chapter 1. Effects of pesticides, temperature and fish species on the physiological stress response – literature review | 1 |
| 1.1 Introduction | 1 |
| 1.2 Physiological Stress Response | 2 |
| 1.2.1 Adrenal Endocrinology - Structure and Function of the Teleostean Head Kidney | 2 |
| 1.2.2 Physiological Effects of Corticosteroid Hormones | 4 |
| 1.3 Pesticides | 7 |
| 1.3.1 Pesticide Uses and Application | 7 |
| 1.3.2 Pesticide Exposure | 8 |
| 1.3.3 Organophosphates and Carbamates | 9 |
| 1.3.4 Effects of Pesticides in Fish | 15 |
| 1.4 Temperature | 17 |
| 1.4.1 Temperature and Projected Effects of Global Climate Change on Aquatic Ecosystems | 17 |
| 1.4.2 Projected Effects of Climate Change on Fish | 19 |
| 1.4.3 Temperature and Pesticides | 21 |
| 1.5 Species Differences | 22 |
| 1.5.1 Species Differences and Stress | 23 |
| 1.5.2 Species Differences and Pesticides | 24 |
| 1.5.3 Species Differences and Temperature | 24 |
| 1.6 Summary | 25 |
| 1.7 References | 27 |
| Chapter 2. The physiological stress response indicators in whitefish (<i>Prosopium williamsoni</i>) and suckers (<i>Catostomus sp.</i>) sampled along a gradient of temperature and agricultural chemicals in the Oldman river, Alberta | 42 |
| 2.1 Introduction | 42 |
| 2.2 Materials and Methods | 45 |
| 2.2.1 Description of the Sites and Fish Species | 45 |
| 2.2.2 Capture and Sampling of the Fish | 48 |
| 2.2.3 Acetylcholinesterase (AChE) | 51 |
| 2.2.4 Physiological Stress Response | 51 |
| 2.2.5 Statistical Analysis | 52 |
| 2.3 Results | 52 |
| 2.3.1 Pesticide Exposure | 52 |
| 2.3.2 Temperature Regimes | 54 |

| | |
|---|-----|
| 2.3.4 Physiological Stress Response | 54 |
| 2.4 Discussion | 56 |
| 2.4.1 Exposure | 56 |
| 2.4.2 Physiological Stress Response | 62 |
| 2.5 Conclusions | 67 |
| 2.6 References | 68 |
| Chapter 3. Effects of Dimethoate and temperature on acetylcholinesterase activity and cortisol secretion in rainbow trout (<i>Oncorhynchus mykiss</i>): <i>in Vivo</i> and <i>in Vitro</i> exposure | 75 |
| 3.1 Introduction..... | 75 |
| 3.2 Materials and Methods..... | 77 |
| 3.2.1 <i>In Vivo</i> Exposure to Waterborne Dimethoate (pure grade)..... | 78 |
| 3.2.2. <i>In Vitro</i> Exposure of Head Kidney Cells to Dimethoate..... | 82 |
| 3.2.3 Statistical Analysis | 84 |
| 3.3 Results..... | 84 |
| 3.3.1 <i>In Vivo</i> Exposure to Dimethoate | 84 |
| 3.3.2 <i>In Vitro</i> Exposure of Head Kidney Cells to Dimethoate (pure grade)..... | 90 |
| 3.3.3 <i>In Vitro</i> Exposure of Head Kidney Cells to Commercial Dimethoate (Cygon) | 93 |
| 3.4 Discussion..... | 96 |
| 3.4.1 AChE Activity – A Marker of Pesticide Exposure | 96 |
| 3.4.2 Physiological Stress Response | 97 |
| 3.5 Conclusions..... | 101 |
| 3.6 References..... | 103 |
| Chapter 4. General Discussion and Conclusion..... | 109 |
| 4.1 General Overview | 109 |
| 4.2 Oldman Study Conclusions..... | 109 |
| 4.3 Dimethoate Study Conclusions..... | 111 |
| 4.4 Concluding Statements | 112 |
| 4.5 References..... | 114 |
| Appendix 1..... | 115 |

List of Tables

| | |
|--|-----|
| Table 2.1. Characteristics of sites sampled in the Oldman river, Alberta. | 47 |
| Table 2.2. Water concentrations of pesticides detected at sites situated along the Oldman River, Alberta..... | 49 |
| Table 2.3. Plasma Glucose (mean±SE, n) in whitefish and suckers sampled at sites along the Oldman river in early summer 2004, late summer 2004, spring 2005, late summer 2005..... | 58 |
| Table 2.4. Condition factor (mean±SE, n) in whitefish and suckers sampled at sites along the Oldman river in early summer 2004, late summer 2004, spring 2005, late summer 2005..... | 59 |
| Table 3.1. Temperature and water chemistry (mean ± SEM) during Dimethoate (pure grade) and temperature treatments..... | 79 |
| Table 3.2. AChE activity inhibition (mean±SE, n) in brain, muscle, and plasma of rainbow trout exposed to Dimethoate..... | 85 |
| Table 3.3. Plasma glucose, liver glycogen, condition factor and liver somatic index, (mean ± SE, n), in rainbow trout exposed to Dimethoate (pure grade) at two different temperatures | 89 |
| Table A.1. AChE activity in Salmonids and Catostomids exposed to OP, CB, or Field pesticides..... | 115 |
| Table A.2. Physiological indicators of stress in fish exposed to pesticides..... | 121 |
| Table A.3. Behavioral changes in fish exposed to OP and CB pesticides..... | 125 |
| Table A.4. Effect of temperature on the stress response in fish | 129 |
| Table A.5. Effect of temperature on pesticide exposure and AChE activity in fish..... | 132 |
| Table A.6. Species differences in AChE activity and the responses to pesticide exposure and temperature..... | 134 |

List of Figures

| | |
|---|----|
| Figure 1.1. The basic structure of organophosphate and carbamate pesticides | 10 |
| Figure 2.1. Location of sampling sites in the Oldman River Basin, Southern Alberta .. | 46 |
| Figure 2.2. Mean water temperatures for each site during sampling periods | 50 |
| Figure 2.3. Plasma AChE (mean±SE) in whitefish and suckers sampled at sites along the Oldman River..... | 53 |
| Figure 2.4. Plasma cortisol (mean±SE) in whitefish and suckers sampled at sites along the Oldman River..... | 55 |
| Figure 2.5. Liver glycogen (mean±SE) in whitefish and suckers sampled at sites along the Oldman River..... | 57 |
| Figure 3.1. Hourly temperature profile for days 11-21 in the experimental tanks | 79 |
| Figure 3.2. The activity of AChE activity in tissues of rainbow trout exposed to Dimethoate (pure grade) at two different temperatures..... | 87 |
| Figure 3.3. A) Plasma cortisol and (B) basal and ACTH-stimulated cortisol secretion (mean±SE) by isolated head kidneys of rainbow trout exposed to Dimethoate (pure grade) at two different temperatures..... | 88 |
| Figure 3.4. Gill Na ⁺ /K ⁺ ATPase activity (mean±SE) in rainbow trout exposed to Dimethoate (pure grade) at two different temperatures..... | 89 |
| Figure 3.5. Cortisol secretion (ACTH-stimulated) and viability of head kidney cells exposed to Dimethoate (pure grade) in vitro at 11°C and 20°C | 92 |
| Figure 3.6. Cortisol secretion (ACTH-stimulated) and viability of head kidney cells exposed to Dimethoate (commercial grade) in vitro at 11°C and 20°C | 95 |

List of Abbreviations

| | |
|---|---|
| ACh | acetylcholine |
| AChE | acetylcholinesterase |
| ACTH | adrenocorticotrophic hormone |
| CB | carbarnates |
| CBG | corticosteroid-binding globulin |
| ChE | cholinesterase |
| CRH | corticotrophin releasing hormone |
| DAG | diacylglycerol |
| EB | ethylene bisdithiocarbamate, |
| ER | endoplasmic reticulum |
| Ft. M | Fort MacLeod |
| HPI | hypothalamo-pituitary-interrenal |
| LDL | low-density lipoproteins |
| Mon | Monarch |
| Na ⁺ /K ⁺ -ATPase | Na ⁺ /K ⁺ -dependent adenosine triphosphatase |
| OC | organochlorine |
| OP | organophosphorus insecticides |
| Pav | Pavan |
| Per | Pearce Road |
| PKA | protein kinase A |
| PKC | protein kinase C |
| Pop | Popson Park |
| PY | pyrethroid |
| StAR protein | Steroidogenic Acute Regulatory protein |
| Sum | Summerview |
| TR | triazine |

Chapter 1.

Effects of pesticides, temperature and fish species on the physiological stress response – literature review

1.1 Introduction

Fish live in a wide range of habitats all over the world. However, many of their natural habitats are being altered by anthropogenic influences, including physical structures (dams and weirs), agricultural runoff, industrial waste, and urban pollutants. Rivers and lakes are also threatened by global climate change which could lead to changes in water temperature. Fish are exothermic and exposures to warmer water temperatures, particularly in cold-water fish species, can significantly alter their physiological status. Invasion by non-native species is another modification of fish habitat, potentially linked to climate change. Any significant alteration to fish habitat, whether it is physical, chemical or biological, can lead to stress, and prolonged or high intensity exposures to stressors have the potential to impact fitness at individual and population level. Different species of fish may respond to specific stressors differently - while some fish may be resistant to certain types of stress, others may be very sensitive. These species-specific responses can disrupt equilibrium that exists within an aquatic system; however our understanding of the various interactions is limited at present. *This literature review will focus on how the physiological stress response is regulated, how pesticides and temperature affect the stress response, and how different fish species respond to these stressors.*

1.2 Physiological Stress Response

1.2.1 Adrenal Endocrinology - Structure and Function of the Teleostean Head Kidney

a. Anatomy

The physiological stress response is mediated by the adrenal system in all vertebrates; however, the structural organization of the adrenals varies. The adrenal gland of mammals is separated into a medulla composed of chromaffin cells secreting catecholamines (epinephrine and norepinephrine) and a cortex composed of adrenocortical cells secreting steroids, including cortisol. In teleosts, the anterior portion of the kidney, known as the head kidney, is analogous to the mammalian adrenal gland (Norris 1997, Mommsen et al. 1999, Hontela 1997, 2005). The head kidney is not separated into a medulla and cortex but there is close physical association between the chromaffin and adrenocortical (also known as interrenal) cells. The head kidney, which has lost its renal function, consists of interspersed islets of adrenocortical cells that are embedded in a matrix of lymphoid tissue (Hontela 2005).

b. Signalling Pathway Leading to Cortisol Secretion

The secretion of the stress hormones corticosteroids is regulated by the hypothalamo-pituitary-interrenal (HPI) axis in teleost fish, as well as other vertebrates. When an organism is exposed to a stressor, within a few seconds the corticotrophin releasing hormone (CRH) is released from the hypothalamus and stimulates the release of adrenocorticotrophic hormone (ACTH) from the pituitary gland (Hontela 2005). The ACTH then stimulates the synthesis and release of corticosteroid hormones, including cortisol, from the adrenals. The biochemical pathways and synthesis of corticosteroids in

teleosts are not as well characterised as those in mammals, but the experiments that have been conducted show a similarity to mammals.

In all vertebrates studied, ACTH targets the adrenocortical cells in the adrenal gland, binds to membrane receptors and stimulates pathways which mobilize cholesterol for production of steroid hormones (Stocco 2000). In mammals, the main pathway involves cAMP and protein kinase A (PKA) while the other involves protein kinase C (PKC) and diacylglycerol (DAG) (Stocco et al. 2005). The importance of ACTH, cAMP, PKA and PKC in the synthesis of cortisol has been also demonstrated in teleost fishes (Patiño et al. 1986, Leblond et al. 2001, Lacroix & Hontela 2001). The cAMP-PKA pathway is activated when ACTH binds to receptors on an adrenocortical membrane and activates G proteins to stimulate adenylate cyclase (Colby & Longhurst 1992, Hontela 1997, Lacroix & Hontela 2001, Stocco et al. 2005). Stimulation of adenylate cyclase increases intracellular levels of cAMP which activates PKA. PKA activation results in the phosphorylation of proteins and transcription factors that are responsible for the activation of the enzyme cholesterol ester hydrolase (Colby & Longhurst 1992, Hontela 1997, Stocco et al. 2005). Cholesterol ester hydrolase releases free cholesterol from cholesterol esters. Cholesterol is supplied to the adrenocortical cells by circulating low-density lipoproteins (LDL) droplets and *de novo* synthesis in the adrenocortical cells (Norris 1997; Stocco 2000). Once cholesterol is present in the outer mitochondrial membrane, it is transported into the inner membrane by the Steroidogenic Acute Regulatory (StAR) protein to be converted into pregnenolone by the side-chain cleavage enzyme, P-450_{scc} (Norris 1997; Geslin & Auperin 2004). Pregnenolone then moves to the smooth endoplasmic reticulum (ER) where it goes through further transformations

(Norris 1997). The conversion of cholesterol into pregnenolone is the rate limiting step in the production of many of the corticosteroids (Colby & Longhurst 1992, Stocco 2000).

c. Cortisol Feedback and Degradation

Secretion of corticosteroids is controlled by a negative feedback loop involving both CRH and ACTH (Colby & Longhurst 1992, Norris 1997, Hontela 1997). When plasma levels of ACTH and corticosteroids are high, they inhibit the release CRH and ACTH, decreasing cortisol synthesis (Colby & Longhurst 1992, Norris 1997, Hontela 1997). The majority of cortisol that is released from the adrenals is bound to a specific plasma protein called corticosteroid-binding globulin (CBG) (Norris 1997, Mommsen et al. 1999, Hontela 2005). Occupied CBG can interact with a target cell membrane which then facilitates dissociation of the glucocorticoid and allows its entrance into the target cell (Norris 1997). The main site for glucocorticoid degradation is in the liver but some degradation may also take place in the kidneys and intestine (Norris 1997, Mommsen et al. 1999).

Glucocorticoids, including cortisol, are metabolized by a series of enzymatic reactions, including transformations by cytochrome P-450 (Norris 1997, Mommsen et al. 1999), increasing water solubility of the steroids and facilitating their excretion via urine, bile or feces.

1.2.2 Physiological Effects of Corticosteroid Hormones

The main corticosteroid in teleosts is cortisol, a steroid that plays a major role in maintaining energy stores and energy mobilization in times of stress (Norris 1997, Hontela 1998, Mommsen et al. 1999, Schreck et al. 2001, Barton 2002). Teleost fish do

not have high levels of mineralocorticoids such as aldosterone, and cortisol has metabolic as well as osmoregulatory effects (Hontela 2005).

a. Metabolic Actions of Cortisol

The main effect of cortisol is energy mobilization by the metabolism of carbohydrates, proteins, amino acids and lipids (Norris 1997). Studies with teleost fish often rely on plasma glucose and liver glycogen stores as indicators of energy metabolism due to stress or exposure to increased cortisol levels. The metabolic actions of cortisol were reviewed by Mommsen et al. (1999). Glucocorticoids are known to increase plasma glucose through degradation and mobilization of proteins, releasing amino acids which are then used through increased activity of gluconeogenic enzymes in hepatic gluconeogenesis. In the liver, cortisol may have some glycogenolytic effects as well; however catecholamines such as epinephrine are mainly responsible for the mobilization of liver glycogen during stress. It has been suggested that during stress, cortisol and insulin interact to not only prevent glycogen breakdown but also increase its synthesis. Thus, as summarized in Mommsen et al. (1999), the overall effect of cortisol in a stressed fish is liver glycogen conservation. Cortisol action on muscle glycogen is not very well understood, but it had been proposed that cortisol may play a role in exercise-induced muscle glycogen mobilization (Milligan 2003). The differences in the stress response and cortisol actions in different species of fish might underlie the variability in responses to cortisol.

b. Osmoregulatory Actions

In freshwater fish cortisol regulates sodium fluxes and stimulates the transfer of sodium across the gill, the mucosa in the gut and the kidney (Norris 1997). The Na^+/K^+ -

dependent adenosine triphosphatase (Na^+/K^+ -ATPase) and Ca^{2+} -ATPase in the gills, gut and kidney is increased when fish are treated with cortisol (Norris 1997). Cortisol is also known to stimulate the branchial calcium pump and increase calcium absorption by the whole body in freshwater rainbow trout, suggesting that cortisol plays a role in fresh water acclimatization (Mommsen et al. 1999). Cortisol is known to increase during the parr-smolt transformation in salmonids, enhancing the hypo-osmoregulatory ability when anadromous fish move into seawater (Mommsen et al. 1999). Cortisol also facilitates the adaptation of salmonids to seawater by stimulating cellular differentiation of chloride cells and branchial and gill Na^+/K^+ -ATPase activity (Mommsen et al. 1999).

c. Reproductive Actions

Cortisol inhibits reproductive functions in both mature and immature fish by affecting production of vitellogenin and oestradiol binding capacity (Mommsen et al. 1999). The stress response increases the amount of cortisol circulating within the body, potentially having a negative effect on the female through decreased gamete quality and progeny fitness (Schreck et al. 2001). Tilapia (*Oreochromis mossambicus*) implanted with cortisol had smaller oocytes, reduced levels of testosterone and 17β -estradiol concentrations (Schreck et al. 2001) and produced smaller larvae (Barton 2002). The effects of these studies clearly show that cortisol has an adverse impact on the reproductive system in many types of fish, and increasingly stressed fish populations may be at a risk of reduced survivorship.

d. Immune Actions

Cortisol production has a negative influence on the immune system in most teleost fish (Barton & Iwama 1991, Hontela 1998, Schreck et al. 2001). The most

extreme example of cortisol's negative impact may be observed in Pacific salmon which all die after spawning. During spawning, Pacific salmon lose control of the pituitary regulatory influences on cortisol secretion by the head kidney and it has been suggested that there is a decrease in the ability to remove cortisol from circulation (Schreck et al. 2001). Cortisol also inhibits the ability of the fish to resist disease or pathogens. High levels of cortisol are known to suppress the ability of fish to form antibodies because the production of leukocytes is depressed (Barton & Iwama 1991, Hontela 1997, Schreck et al. 2001). Studies in brown trout (*Salmo trutta*) have shown that handling stress can cause a significant decrease in leukocytes (Barton & Iwama 1991). When fish are removed from the stressful situation that is causing elevated cortisol levels, the immune system can return to normal (Barton & Iwama 1991, Schreck et al. 2001).

Cortisol also interacts with other hormones, including thyroid hormones, growth hormone and catecholamines (Hontela 1997), particularly during exposures to stressors. The remainder of this review will examine stress with emphasis on cortisol in relation to pesticides, temperature, and species differences. *The effects of multiple stressors (specifically pesticides and temperature) on the physiological stress response and how these stressors affect different species of fish, have not been investigated. These interactions will be the main focus of this thesis.*

1.3 Pesticides

1.3.1 Pesticide Uses and Application

Pesticides are chemicals used to control pests; their primary use is to increase agricultural yields and prevent the spread of vector-borne disease that have implications

to health and the economy (Hayes 1991). Pesticides do not include antibiotics. Pesticides can be classified based on the pest that they are targeting, and include algicides, fungicides, herbicides, insecticides, molluscicides, nematicides, and rodenticides. They are also categorized by their chemical structure; the most commonly used pesticides are organophosphates, carbamates, organochlorines, pyrethroids, triazines, ureas, and aryloxyalkanoic acids (Stenerson 2004).

Pesticides are generally applied during the growing season, but the application times depend on weather, the targeted pest and local development of the pest (Frank & Sirons 1979, Trim & Marcus 1990, Fairchild et al. 1999). Pesticides in Southern Alberta have been detected in the air from April until October (Tuduri et al. 2006b). Waite et al. (1995) found the highest deposition amounts of 2, 4D in mid-June and highest deposition amounts of Triallate in mid-May. Traces of pesticides found outside application areas (e.g. watershed) are usually at the highest concentrations during planting season (Frank & Sirons 1979, Tuduri et al. 2006b). Pesticides are often detected in water due to direct application in water, storm runoff, irrigation tail-water releases, and spray drift or vapour transport (Raupach et al. 2001a, b, Tuduri et al. 2006a, b).

1.3.2 Pesticide Exposure

The main routes of exposure to pesticides are through the integument, ingestion and inhalation, or in fish, uptake through gills (Brooks 1976). Depending on the physical properties of the pesticide, it may enter the blood stream and be carried in solution, bind to proteins, or dissolve in lipid particles (Brooks 1976). Distribution of pesticides and their metabolites depends on their ability to cross cell membranes (Brooks 1976). Many

pesticides biotransform by chemical conversion or enzymatic biotransformation; transformation may increase or decrease their toxicity (Brooks 1976). Original or converted pesticides may need to undergo further transformations into water-soluble conjugates in order to be excreted (Brooks 1976).


Although the use of pesticides has great societal benefits, there are many problems associated with them. Pesticides are biologically active chemicals that pose potential harm to non-target organisms (Hayes 1991). People who are involved with the manufacture or use of pesticides are at high risk of exposure (Marrs 1993). Pesticides can also affect domestic and wild animals that are exposed either directly to the pesticide, its runoff, or by eating contaminated food (Brooks 1976, Hayes 1991). Acute pesticide exposure due to runoff caused reductions in fish populations and large scale fish kills (Trim & Marcus 1990, Fairchild et al. 1999, Fulton et al. 1999). Fairchild et al. (1999) determined that the pesticide Matacil 1.8 was responsible for population reductions of Atlantic salmon in New Brunswick and Newfoundland. Chronic exposures to small amounts of pesticides are known to increase incidence of disease, increase stress, and cause behavioural changes in fish (Austin 1999). *This review will focus on OPs and CBs, and their chemical make-up, and mode of action.*

1.3.3. Organophosphates and Carbamates

a. History and Uses


Organophosphorus insecticides (OPs) were developed in Germany during the 1930s as potential chemical warfare agents (Fulton & Key 2001). The OPs together with carbamates (CBs) (Figure 1.1) have become the most widely used classes of insecticides

Organophosphate Esters



X alkyl
alkoxy
Y amido
Z aryl
alkyl
alkoxy

Carbamate Esters



R aryl
alkyl

Figure 1.1. The basic structure of organophosphate and carbamate pesticides.

worldwide (Fulton & Key 2001). In 1998, there were 15 types of carbamates and 34 types of OPs sold in Alberta (Byrtus 2000). They are used to control insects on fruits, vegetables, grain crops, and stored seeds, and in the household, to control cockroaches, houseflies, termites, and protect plants of horticultural interest (Fulton & Key 2001). The OPs and CBs have replaced many organochlorine pesticides because they are less persistent in the environment and biomagnification is not a major issue (Tuduri et al. 2006b). A few problems associated with OPs and CBs are their lack of specificity and high water solubility; they are able to enter the aquatic environment where they are toxic to both vertebrates and invertebrates (Tuduri et al. 2006b). Furthermore, under specific conditions, some OPs remain active in soils for up to six months after application, thus increasing the duration they may be harmful to non-target species (Köprücü et al. 2006). OPs have several mechanisms of degradation in natural waters such as oxidation, photodegradation, and hydrolysis, described in a review by Pehkonen & Zhang (2000).

b. ACh and AChE Function in the Nervous System

The OPs and CBs act through the nervous system, inhibiting cholinesterase (ChE) enzymes (Stenerson 2004). Communication between neurons and nerve, muscle or endocrine cells is chemically mediated - an impulse causes the release of the acetylcholine (ACh) from the axon terminal of a presynaptic neuron into the synaptic cleft (Shankland 1976, Stenerson 2004). ACh binds to a receptor on the postsynaptic membrane and the chemical signal is converted into an electrical signal (Shankland 1976). The electrical signal continues until ACh is removed from the receptor on the postsynaptic membrane by acetylcholinesterase (AChE). AChE hydrolyses ACh into acetate and choline, thus stopping the electrical signal to the postsynaptic membrane

(O'Brien 1976, Beauvais et al. 2001). AChE hydrolyzes ACh by forming an enzyme-substrate complex where ACh acetylates AChE and releases choline (O'Brain 1976, Stenerson 2004). AChE is then hydrolyzed and restored (Stenerson 2004); this reaction occurs very fast (O'Brain 1976, Stenerson 2004). There are two main types of cholinesterase enzymes: acetylcholinesterase (true) and butyrylcholinesterase (pseudo). *This review and thesis will focus on acetylcholinesterase.*

c. OP/CB Mode of Action – AChE Inhibition

OPs and CBs inhibit AChE by targeting the serine hydroxyl group on the AChE active site where they bind to and inactivate the enzyme (Grue et al. 1991, Jett et al. 1993, Abbas & Hayton 1997, Stenerson 2004). OPs form a complex with AChE similar to the enzyme-substrate complex made with ACh, but OPs phosphorylate the serine hydroxyl group (O'Brain 1976, Marrs 1993). The reactivation of AChE by dephosphorylation is very slow (considered irreversible) and controls the overall rate of the reaction (O'Brain 1976, Marrs 1993). CBs act differently than OPs by carbamylating the serine hydroxyl group of AChE. The reactivation by decarbamylation step occurs slowly but is not considered irreversible (O'Brain 1976). When AChE is phosphorylated or carbamylated, it is prevented from hydrolyzing ACh which accumulates in the synaptic cleft (Jett et al. 1993, Marrs 1993, Stenerson 2004). The activation of receptors is thus prolonged, leading to an overstimulation of the postsynaptic cholinergic receptors (Jett et al. 1993, Liu & Pope 1996, Jones et al. 1998) and death.

d. Recovery and Adaptations to OP/CB Exposure

Recovery of AChE inhibition by OPs can occur by dephosphorylation or synthesis of new AChE, which are both slow processes (Grue et al. 1991). Recovery of

brain AChE inhibition in mosquitofish (*Gambusia affinis*) exposed to OPs occurred between 45 and 60 days post exposure, however, muscle AChE only partially recovered and there was no recovery of liver AChE (Carr et al. 1997). There is evidence that some organisms develop adaptations to AChE inhibition (Beauvais et al. 2001). A study by Abbas and Hayton (1997) reported that rainbow trout can increase the synthesis of AChE by 10%. The downregulation of cholinergic receptors was found to be an adaptive response to overstimulation of the postsynaptic cell (Jett et al. 1993, Jones et al. 1998). Rainbow trout exposed to carbaryl and permethrin decreased the number of cholinergic receptors compared to controls (Jones et al. 1998). Liu and Pope (1996) demonstrated that rats exposed to chlorpyrifos decreased ACh synthesis which reduced the number of ACh molecules present in the synapse, thus lessening the effect of the AChE inhibition. Aliesterases and carboxylesterases in some species of fish are suggested to be a protective mechanism because they bind to the active metabolite of some insecticides, thus preventing OPs and CBs from binding to AChE (Clement 1984, Carr et al. 1997).

e. OP/CB Mechanisms for Toxicity

Many OPs need to be metabolized to an oxon form while carbamates and some OPs inhibit AChE directly, without biotransformation (Grue et al. 1991, Hoffmann & Papendorf 2006). Habig et al. (1988) found that activation of the oxygen analog was required for malathion and parathion to inhibit AChE in channel catfish (*Ictalurus punctatus*). It should be noted, however, that some CB pesticides that act as fungicides (e.g. IPBC and Mancozeb) have been reported to increase AChE rather than inhibit AChE through unknown mechanisms (Jarrard et al. 2004). Areas of high agricultural intensity may use many different kinds of pesticides, some in small amounts. However,

even exposure to trace amounts of different OPs and CBs can cause AChE inhibition because most OPs and CBs have the same mode of action and have additive effects (Grue et al. 1991; Scholz et al. 2006). It has been proposed that in order for an OP or CB to cause toxicity four processes must occur: 1) binding and inhibition of an extensive number of AChE molecules by the OP or CB, causing impairment of ACh degradation, 2) accumulation of ACh in the synaptic cleft of central and/or peripheral neurons, 3) the excessive and continuous stimulation of postsynaptic receptors, and 4) an altered cellular response due to overstimulation of postsynaptic receptors (Pope 1999).

f. Effects of AChE Inhibition in fish

AChE in most vertebrates is present in the nervous tissue, including neuromuscular junctions (O'Brian 1976) and also in some non-nervous tissue such as erythrocytes (Stenerson 2004). The function of AChE in erythrocytes is unknown but it allows for a simple and non lethal way of detecting exposure to OPs and carbamates (Stenerson 2004). OPs and CBs can significantly affect AChE activity in fish, as documented in Table A.1 (Appendix). In some fish such as *Pocilia reticulata*, brain AChE activity can be inhibited by 70% without causing mortality in acute exposures (Frasco & Guilhermino 2002), while in estuarine fish, inhibition > 70% was associated with mortality and sublethal effects on stamina were detected with inhibitions as low as 50% (Fulton & Key 2001). Traces of OPs and CBs have also been found in the tissues of exposed fish. Sapozhnikova et al. (2004) found detectable amounts of OPs in the liver, gonads, gills and muscle of fish in the Salton Sea, California. OPs and CBs are not distributed in tissues equally; Abbas and Hayton (1997) reported that concentrations of paraxon were higher in the liver followed by the brain, heart, and kidney. Moreover,

even though the liver had the highest amount of paraxon in the tissue, the brain was more sensitive to AChE inhibition.

A negative relationship between brain size and AChE activity has been demonstrated in bluegill (*Lepomis macrochirus*) (Beauvais et al. 2002) and golden shiner (*Notemigonus crysoleucas*) but not in rainbow trout (Phillips et al. 2002). Body length and weight had a negative correlation to AChE activity in three-spined stickleback (*Gasterosteus aculeatus*) (Strum et al. 1999) and rainbow trout (Zinkl et al. 1987), whereas Sandahl and Jenkins (2002) did not find a relationship between weight and AChE activity in juvenile steelhead (*Oncorhynchus mykiss*).

1.3.4 Effects of Pesticide in Fish

a. Effects on the Stress Response

Pesticides are toxicants that can activate the stress response in fish, but some can also inhibit cortisol secretion by disrupting the signalling pathways leading to cortisol at specific steps before or after the generation of cAMP (Leblond & Hontela 1999, Leblond et al. 2001, Bisson & Hontela 2002). Table A.2 shows how different pesticides affect the physiological stress response. The understanding of the effects of pesticides on the stress response is limited at present. *This thesis will provide new data on the effects of OP pesticide exposure and how it affects the physiological stress response, under both laboratory and field conditions.*

b. Effects on Behaviour and Development

Sublethal inhibition of AChE activity affects various behavioural endpoints in fish, including swimming capacity and activity (Little & Finger 1990, Beauvais et al.

2001, Table A.3). Changes in behaviour can have consequences to fish by reducing feeding, as well as causing problems avoiding predators, finding mates and migrating (Dutta et al. 1992, Beauvais et al. 2001). OP and CB exposure can affect reproduction and reproductive potential (Bhattacharya 1993, Begum & Vijayaraghavan 1995). Many OPs and CBs cause developmental problems in embryos and juveniles; diazinon caused deformities in larval medaka (*Oryzias latipes*), including growth retardation and formation of edema in yolk sac (Hamm et al. 1998). Chlorpyrifos, another OP, caused malformations (tail abnormalities, ventral swelling, fin damage, haemorrhaging) and morbidity in juvenile guppies, *Poecilia reticulata*, (De Silva & Samayawardhena 2002).

c. Effects on Food Sources

OPs and CBs in water can further affect fish by altering the food webs, including effects on algae and freshwater phytoplankton (DeLorenzo et al. 2001, Ratageri et al. 2006, Sweilum 2006). Dimethoate decreased levels of algal protein and carbohydrates, inhibiting growth and decreasing cell density in *Scenedesmus incrassatulus* at very low concentrations (0.075 mg/L) (DeLorenzo et al. 2001). Dimethoate decreased primary and gross productivity in a water body by 70% after 12 hrs of exposure (Ratageri et al. 2006). Studies in lentic systems exposed to the OP phosphamidon, reported significant reductions in both phytoplankton and zooplankton (Dhar et al. 2004). Pesticides are very useful for protection against pests; however, they can be detrimental to non-target species. As pesticide use continues to increase (Bloomfield et al. 2006), it is important to better understand how pesticides affect non-target species and what the fates of pesticides are in the environment, specifically as the climate changes and water temperatures increase.

1.4 Temperature

1.4.1 Temperature and Projected Effects of Global Climate Change on Aquatic Ecosystems

Human activities such as deforestation, cultivation, irrigation, dam construction and industrial operations can have a negative impact on water quality; moreover, some of these activities lead to an increase in water temperature (Ihnat & Bulkley 1984, Beitinger et al. 2000). Global climate change and specifically an increase in global temperature is also of great concern (De Stasio et al. 1996, Eaton & Scheller 1996, Fang et al. 2004a, b) since with the change in global temperatures an increase in both surface and ground-water temperatures is projected. The Experimental Lakes area in Ontario has already seen a 2°C increase in water temperature in the last twenty years (Morgan et al. 1998). Using modeling and the scenario of doubling of atmospheric carbon dioxide, an increase in mean water temperatures by 1-7°C has been predicted for many lakes and rivers (De Stasio et al. 1996). Studies on the Fraser River, British Columbia, Canada predict water temperature increases of 0.022°C per year (Morrison et al. 2002).

Increases in water temperature are projected to have significant impacts on fish habitats and fish populations, however, the impacts depend on the type of fish (whether it belongs to a cold-, cool- or warm-water fish guild) and the type of aquatic system it inhabits. Fish have rather precise thermal requirements and often confine their distributions to specific water temperatures (Brandt 1993). Body temperatures of most fish are a direct function of water temperature and a combination of environmental, anatomical and physiological factors cause the body temperature of most fish to be

usually only 0.1 to 1°C above ambient water temperatures (Beitinger et al. 2000). Temperature can be viewed as an environmental resource in which fish will compete with each other for favourable temperatures (Beitinger et al. 2000, Beitinger & Magnuson 1975). Field studies have shown that most fish have a total distribution range limited to a 10°C window of their thermal preference and will spend most of their time within 2°C around their thermal preference (Brandt 1993).

The optimal temperature for fish is important because it is the temperature where fish can feed optimally, grow, and complete their life cycle (Brandt 1993, Kroger et al. 1999, Fang et al. 2000a, Myrick & Cech 2004). Fish may be able to live in water temperatures above or below their optimal temperature but are then usually not fit enough to pass their genes on to the next generation (Eaton & Scheller 1996). A review by Beitinger et al. (2000) concluded that the family Salmonidae had the lowest tolerance to high temperatures, as compared to many other fish families.

Temperature can affect not only the physiology but also the habitat of the fish. Changes in stratification within lakes and river pools and decreasing dissolved oxygen will force fish out of their existing habitats and increase competition within and between species for more suitable habitat (Stefan et al. 1993, De Stasio et al. 1996, Fang et al. 2004a, b, Tate & Lancaster 2007). The effects of increasing temperatures and changes in climate patterns are important to fish well being as fish habitat and life histories revolve around water temperatures (De Stasio et al. 1996, Eaton & Scheller 1996, Rahel et al. 1996, Fang et al. 2004a, b).

1.4.2 Projected Effects of Climate Change on Fish

Climate change is predicted to affect both the number of fish kills associated with temperature and increase the number of fish invasions (Trim & Marcus 1990). Fish kills are predicted to increase in both summer and winter: in summer through excessive water temperatures, low dissolved oxygen and increased salinity while in winter only through milder temperatures causing low dissolved oxygen (Trim & Marcus 1990, Fang et al. 2000 a, b, Sapozhnikova et al. 2004). As water temperatures increase, warm water fish may invade lakes and rivers they were previously unable to inhabit due to the low water temperature (Eaton & Scheller 1996, Morgan et al. 1998). These invasions could further exacerbate the effect that global climate change has on a particular ecosystem.

There is the possibility, however, that if raising water temperatures is a slow process, cool- and cold-water fish may be able to acclimatize to warmer waters. Acclimatisation temperature is suggested to be the most important factor influencing temperature tolerance (Cincotta & Stauffer 1984, Beitinger & Bennett 2000, Beitinger et al. 2000). Studies have shown that fish are able to increase their tolerance of high temperatures by more than 10°C when acclimatised to a warm temperature (Cincotta & Stauffer 1984, Ihnat & Bulkley 1984, Beitinger et al. 2000). For example, upper avoidance temperatures in white suckers (*Catostomus commersoni*) changed from 15°C when acclimatised to 6°C, to 30°C when acclimatised at 24°C (Cincotta & Stauffer 1984). Another study in mountain whitefish (*Prosopium williamsoni*) found that fish acclimatised to 15°C preferred warmer temperatures than fish acclimatised to 5°C (Ihnat & Bulkley 1984).

Non-lethal temperature increases can have enormous impacts on fish communities because thermal stress causes fish to divert energy from growth to homeostasis maintenance, including the induction of heat-shock proteins (Iwama et al. 1998, Viant et al. 2003). Both increases and decreases in water temperature can cause reductions in feeding and growth rates (Brett 1971, Bartell et al. 1986, Brandt 1993). Low water temperatures are known to decrease reproduction and fecundity in medaka (Koger et al. 1999). In mountain whitefish, high water temperatures during egg development result in increased egg mortality: 100% mortality at 15°C and 12°C, 97.8 to 100% mortality at 11°C, 34.4 to 95.2% mortality at 10°C, and 6.5 to 9.5% mortality at 6°C (Rajagopal 1979).

Exposure to water temperatures outside the thermal preference of a fish is known to activate the physiological stress response (Table A.4), demonstrated through endpoints such as an increase in cortisol levels (Barton & Peter 1982, Davis & Parker 1990, Davis 2004, Weber & Bosworth 2005, Davis & Peterson 2006), decrease or increase in plasma glucose (Davis & Parker 1990, Davis 2004, Lerman et al. 2004) and decrease in liver glycogen (Viant et al. 2003, Lerman et al. 2004). More information is needed to determine how fish will respond to temperatures changes. This thesis will evaluate how the physiological stress response in fish is affected by temperature. However, it is hypothesised that how a fish responds to temperature depends on the degree of temperature change, acclimatisation temperature, and species of fish.

Temperature also plays an important role in life history characteristics - a study of white suckers reported that fish living in colder subpolar regions took longer to reach maturity and had slower growth rates than fish in warmer temperate regions (Duchesne &

Magnan 1997). Behaviour can be influenced by non-thermal stimuli which can override the preferred temperature response (Beitinger & Magnuson 1975). Juvenile bluegill preferred 31°C over 27° or 34°C but when adult bluegill were present, the juveniles spent the majority of their time outside of the preferred 31°C. Temperature can also affect how a fish will respond to toxicants, including pesticides, in the environment.

1.4.3 Temperature and Pesticides

Global climate change is predicted to have far reaching effects on agriculture. Predicted changes in weather patterns will force changes in farming practices and the use of chemicals associated with agriculture. Canada and the United Kingdom are projected to have milder, wetter winters and hotter drier summers, with an overall change in average temperature and a longer growing season (Smith & Almaraz 2004, Bloomfield et al. 2006). Changes in temperature and growing season affect agricultural production, types of crops that can be grown and the pests associated with the introduction of new crops (Smith & Almaraz 2004, Bloomfield et al. 2006). Changes in crop type and invasion of new pests will shift pesticide use as well (Smith & Almaraz 2004). Climate change is predicted to increase weeds, pests and diseases, which will most likely lead to an increase in pesticide use (Bloomfield et al. 2006).

Changes in temperature are also predicted to change how pesticides act in non-target species such as fish. Temperature is predicted to change pesticide exposures by increasing rates of dispersal and increasing transport distances from source, however, temperature will also increase degradation rates (Bloomfield et al. 2006). The effect of temperature on the toxicity of a pesticide depends on the pesticide itself: some pesticides

are more toxic in warm water while others are more toxic in cold water (Capkin et al. 2006). In fish, the mode of action of OPs and CBs can be temperature-dependent, making the fish more susceptible to OPs and CBs at higher temperatures due to temperature effects on metabolic rate, availability of oxygen in tissues and waste production (Macek et al. 1969). There are conflicting studies on the effect that temperature has on AChE. Studies by Beauvais et al. (2002) in bluegill, Hazel (1969) in mummichog (*Fundulus heteroclitus*), and Phillips et al. (2002) in walleye (*Stizostedion vitreum*) concluded that temperature does not affect AChE, whereas Hogan (1970) in bluegill, Zinkl et al. (1987) in rainbow trout and Hazel (1969) in goldfish (*Carassius auratus*) showed that AChE increases with increasing water temperature. Activity of AChE may not be significantly different if the temperatures are above 10°C and span less than 12°C (Beauvais et al. 2002). A summary of the effects of temperature on fish is presented in Table A.5. *This thesis will investigate the effects of pesticides on the physiological stress response in fish and the interactions with temperature.*

1.5 Species Differences

Different species respond to adverse conditions in different manners. Species-specific differences have been found in regards to toxicity of pathogens (Gordon et al. 2006), AChE inhibition (Davies et al. 1994, Fulton & Key 2001), metal distribution and metallothionein concentration (Linde et al. 1999), stress and the stress response (Lacroix & Hontela 2004, Jentoft et al. 2005), and vulnerability to oxidative stress (Hasspieler et al. 1994). An extensive review was conducted by Vittozzi and De Angelis (1991) comparing the LC₅₀ of common test species to a wide range of toxicants. Table A.6

provides a summary of how different species respond to stress, pesticides and temperature. *Very few studies examined species differences in fish that share the same environment; this thesis (Chapter 2) will examine how different species living in the same environment respond to stress, pesticide exposure and temperature.*

1.5.1 Species Differences and Stress

Species differ in their ability to respond to stressors. This ability depends on life history, strain and acclimation temperature (Jentoft et al. 2005). Differences in the stress response have been found in domesticated versus non-domesticated fish (Barton 2000, Jentoft et al. 2005). For example, domesticated rainbow trout was able to habituate to stress while non-domesticated Eurasian perch (*Perca fluviatilis*) were not (Jentoft et al. 2005). When different species are similarly stressed, they may have different capacities to respond to this stress, as indicated by variations in corticosteroid-binding proteins, the metabolic clearance rate of cortisol (Barton 2000) and carbohydrate utilization (Jentoft et al. 2005). When lake trout (*Salvelinus namaycush*) and rainbow trout were exposed to a 30 second handling stress, lake trout had higher levels of plasma cortisol while rainbow trout had higher levels of plasma glucose (Barton 2000). Knowing how fish will respond to stressors is important for aquaculture, and for predicting how fish will respond to environmental challenges, including chemical pollutants and identifying species at risk (Barton 2000).

1.5.2 Species Differences and Pesticides

Pesticides can have different effects on different species (Johnson & Wallace 1987). Sapozhnikova et al. (2004) reported that tilapia exposed in the laboratory had more dimethoate and *pp*-DDE in tissues than covina (*Cynoscion xanthulu*), but covina had more malathion, endrin, and DDD concentrations in tissues as compared to tilapia. Inhibition of AChE activity can be different between species; furthermore the inhibition of different tissues can also be different between species (Carr et al. 1997). One study found that brain tissue in mosquitofish had greater inhibition of AChE activity than largemouth bass (*Micropterus salmoides*), bluegill sunfish and golden shiners but mosquitofish and golden shiners had lower inhibition of muscle AChE activity compared to largemouth bass and bluegill (Carr et al. 1997). Specific activity of AChE can also be different between species; Johnson and Wallace (1987) and Chuiko (2000) found that control levels of AChE activity can be up to 7-fold higher in perch (Percidae) compared to whitefish (Coregonidae). The differences in species reaction to pesticide exposure and AChE inhibition could be due to different levels of aliesterases which have been suggested to provide a protective mechanism to OP exposure (Carr et al. 1997). Species differences to pesticide exposure may be caused by differences in uptake and metabolism of the pesticide (Carr et al. 1997), affinity of the pesticide for AChE (Keizer et al. 1995), amount of enzyme, the rate of AChE phosphorylation (Johnson & Wallace 1987, Intorre et al. 2004) and kinetics of AChE inactivation (Johnson & Wallace 1987).

1.5.3 Species Differences and Temperature

It is well known that different species of fish prefer different water temperatures - fish are even classified according to their thermal preference: cold-water fish have a maximum temperature tolerance under 24.3°C, cool water fish have a maximum temperature tolerance between 26.5 and 29.9°C, and warm water fish have a maximum temperature tolerance over 30°C (Eaton & Scheller 1996). Temperature differences between similar species can be caused by differences in fish size, as metabolic rates decreases when fish are larger (Galbreath et al. 2004). Metabolic rates in polar fish tend to be higher at low temperatures making them better suited for cold water than non-polar fish (van Dijk et al. 1999). High water temperatures cause fish to accumulate anaerobic end products due to an insufficient supply of oxygen to the tissues (van Dijk et al. 1999). This causes a failure in the electron transport chain and will eventually lead to death in some fish, while fish that can handle warm water either clear out the anaerobic end products or have enough oxygen available to the tissues to prevent build-up of anaerobic end products (van Dijk et al. 1999). Temperature is an important consideration in aquaculture - knowing how different species of fish will respond to cold shock, heat shock, and warming of water can influence where aquaculture activities can take place and how a rise in water temperature will affect the fish (Kuo & Hsieh 2006).

1.6 Summary

Functional integrity of the adrenal gland and the head kidney in teleost fish is very important for the well being of the organism. Excessive stress caused by environmental stressors such as temperature and/or toxicants such as pesticides can have severe consequences to the health and well-being of fish. Furthermore toxicants, such as

pesticides can inhibit normal endocrine function. The adrenal tissue can lose the ability to produce hormones such as cortisol which help fish to properly cope with stressful situations. Excessive stress or an impaired stress response can lead to decreased reproduction, growth, fitness and survivorship. Organisms are often faced with more than one stressor and/or toxicant and it is important to understand how these potentially complex interactions of multiple stressors will affect organisms. The effects of multiple stressors on the physiological stress response are a new area of research and this thesis will examine the stress response in fish exposed to pesticides and warm water temperature. There is very little information regarding how temperature and pesticides affect the stress response and other physiological functions such as acetylcholinesterase activity. Species differences can account for some of the variation in the literature about temperature and pesticide stress but further research is needed. Furthermore, due to the continued use and exposure of organisms to toxicants such as pesticides and to changes in global climate, more research is needed to determine how fish respond to these stressors both in the laboratory and in the field.

The overall objective of this study is to determine, through the use of physiological markers (plasma cortisol, liver glycogen, plasma glucose, condition factor, AChE activity), how temperature and pesticides affect fish. Furthermore, this study will examine how different species of fish that live in the same environment will respond to stressors. The interactive effects of pesticide exposure and temperature on the physiological status will be quantified, and species-specific differences in vulnerability will be assessed, providing new information on the physiological stress response to multiple stressors in fish.

1.7 References

- Abbas, R. & Hayton, W.L. 1997. A physiologically based pharmacokinetic and pharmacodynamic model for paraoxon in rainbow trout. *Toxicology and Applied Pharmacology* 145:192-201.
- Altinok, I., Capkin, E., Karahan, S., & Boran, M. 2006. Effects of water quality and fish size on toxicity of methiocarb, a carbamate pesticide, to rainbow trout. *Environmental Toxicology and Pharmacology* 22:20-26.
- Alvarez, M.d.C. & Fuiman, L.A. 2006. Ecological performance of red drum (*Sciaenops ocellatus*) larvae exposed to environmental levels of the insecticide malathion. *Environmental Toxicology and Chemistry* 25:1426-1432.
- Austin, B. 1999. The effects of pollution on fish health. *Journal of Applied Microbiology Symposium Supplement* 85:234S-242S.
- Bartell, S.M., Breck, J.E., Gardner, R.H., & Brenkert, A.L. 1986. Individual parameter perturbation and error analysis of fish bioenergetics models. *Canadian Journal of Fisheries and Aquatic Sciences* 43:160-168.
- Barton, B.A. 2000. Salmonid fishes differ in their cortisol and glucose responses to handling and transport stress. *North American Journal of Aquaculture* 62:12-18.
- Barton, B.A. 2002. Stress in fishes: A diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and Comparative Biology* 42:517-525.
- Barton, B.A. & Iwama, G.K. 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases* 1:3-26.
- Barton, B.A. & Peter, R.E. 1982. Plasma cortisol stress response in fingerling rainbow trout *Salmo gairdneri* Richardson, to various transport conditions, anaesthesia, and cold shock. *Journal of Fish Biology* 20:39-51.
- Barton, B.A. & Schreck, C.B. 1987. Influence of acclimation temperature on interrenal and carbohydrate stress responses in juvenile Chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 62:299-310.
- Beauvais, S.L., Cole, K.J., Atchison, G.J., & Coffey, M. 2002. Factors affecting brain cholinesterase activity in bluegill (*Lepomis macrochirus*). *Water, Air, and Soil Pollution* 135:249-264.
- Beauvais, S.L. Jones, S.B., Brewer, S.K., & Little, E.E. 2000. Physiological measures of neurotoxicity of diazinon and malathion to larval rainbow trout (*Oncorhynchus*

- mykiss*) and their correlation with behavioural measures. *Environmental Toxicology and Chemistry* 19:1875-1880.
- Beauvais, S.L., Jones, S.B., Parris, J.T., Brewer, S.K., & Little, E.E. 2001. Cholinergic and behavioural neurotoxicity of carbaryl and cadmium to larval rainbow trout (*Oncorhynchus mykiss*). *Ecotoxicology and Environmental Safety* 49:84-90.
- Begum, G. 2004. Carbofuran insecticide induced biochemical alterations in liver and muscle tissues of the fish *Clarias batrachus* (linn) and recovery response. *Aquatic Toxicology* 66:83-92.
- Begum, G. & Vijayaraghavan, S. 1995. Chronic effects of dimethoate on the reproductive potential of the fresh-water teleost, *Clarias batrachus*. *Pesticide Science* 44:233-236.
- Begum, G. & Vijayaraghavan, S. 1999. Effect of acute exposure of the organophosphate insecticide Rogor on some biochemical aspects of *Clarias batrachus* (Linnaeus). *Environmental Research* 80A:80-83.
- Beitinger, T.L. & Bennett, W.A. 2000. Quantification of the role of acclimation temperature in temperature tolerance of fishes. *Environmental Biology of Fishes* 58:277-288.
- Beitinger, T.L., Bennett, W.A., & McCauley, R.W. 2000. Temperature tolerances of North American freshwater fishes exposed to dynamic changes in temperature. *Environmental Biology of Fishes* 58:237-275.
- Beitinger, T.L. & Magnuson, J.J. 1975. Influence of social rank and size on thermoselection behavior of bluegill (*Lepomis macrochirus*). *Journal of the Fisheries Research Board of Canada* 32:2133-2136.
- Bhattacharya, S. 1993. Target and non-target effects of anticholinesterase pesticides in fish in: *Proceeding of the second European conference on toxicology. Science of the Total Environment Supplement*:859-866.
- Bisson, M. & Hontela, A. 2002. Cytotoxic and endocrine-disrupting potential of atrazine, diazinon, endosulfan, and mancozeb in adrenocortical steroidogenic cells of rainbow trout exposed *in Vitro*. *Toxicology and Applied Pharmacology* 180:110-117.
- Bloomfield, J.P., Williams, R.J., Goody, D.C., Cape, J.N., & Guha, P. 2006. Impacts of climate change on the fate and behaviour of pesticides in surface and groundwater – a UK perspective. *Science of the total Environment* 369:163-177.
- Brandt, S.B. 1993. The effect of thermal fronts on fish growth: A bioenergetics evaluation of food and temperature. *Estuaries* 16:142-159.

- Brethead, S., Saglio, P., Saligaut, C., & Auperin, B. 2002. Biochemical and behavioral effects of carbofuran in goldfish (*Carassius auratus*). *Environmental Toxicology and Chemistry* 21:175-181.
- Brett, J.R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). *American Zoologist* 11:99-113.
- Brewer, S.K., Little, E.E., DeLonay, A.J. Beauvais, S.L., Jones, S.B., & Ellersieck, M.R. 2001. Behavioral dysfunctions correlate to altered physiology in rainbow trout (*Oncorhynchus mykiss*) exposed to cholinesterase-inhibiting chemicals. *Archives of Environmental Contamination and Toxicology* 40:70-76.
- Brooks, G.T. 1976. Penetration and distribution of insecticides in: *Insecticide biochemistry and physiology* ed. Wilkinson, C.F. Plenum Press New York 3-58.
- Byrtus, G. 2000. 1998 overview of pesticide sales in Alberta. Available on request at: env.infocent@gov.ab.ca.
- Capkin, E., Altinok, I., & Karahan, S. 2006. Water quality and fish size affect toxicity of endosulfan, an organochlorine pesticide, to rainbow trout. *Chemosphere* 64:1793-1800.
- Carmichael, G.J., Tomasso, J.R., Simco, B.A., & Davis, K.B. 1984. Confinement and water quality-induced stress in largemouth bass. *Transactions of the American Fisheries Society* 113:767-777.
- Carr, R.L., Ho, L.L., & Chambers, J.E. 1997. Selective toxicity of chlorpyrifos to several species of fish during an environmental exposure: Biochemical mechanisms. *Environmental Toxicology and Chemistry* 16: 2369-2374.
- Chuiko, G.M. 2000. Comparative study of acetylcholinesterase and butyrylcholinesterase in brain and serum of several freshwater fish: specific activities and in vitro inhibition by DDVP, an organophosphorus pesticide. *Comparative Biochemistry and Physiology* 127C:233-242.
- Cincotta, D.A. & Stauffer, J.R., Jr. 1984. Temperature preference and avoidance studies of six North American freshwater fish. *Hydrobiologia* 109:173-177.
- Clement, J.G. 1984. Role of aliesterase in organophosphate poisoning. *Fundamental and Applied Toxicology* 4S:96-105
- Colby, H.D. & Longhurst, P.A. 1992. Toxicology of the adrenal gland. In: *Endocrine Toxicology*, edited by C.K. Atterwill and J.D. Flack, Cambridge, UK. Cambridge University Press, pp. 243-281.

- Datta, M. & Kaviraj, A. 2003. Ascorbic acid supplementation of diet for reduction of deltamethrin induced stress in freshwater catfish *Clarias gariepinus*. *Chemosphere* 53:883-888.
- Davies, P.E., Cook, L.S.J., & Goenarso, D. 1994. Sublethal responses to pesticides of several species of Australian freshwater fish and crustaceans and rainbow trout. *Environmental Toxicology and Chemistry* 13:1341-1354.
- Davis, K.B. 2004. Temperature affects physiological stress responses to acute confinement in sunshine bass (*Morone chrysops* X *Morone saxatilis*). *Comparative Biochemistry and Physiology* 139A:433-440.
- Davis, K.B. & Parker, N.C. 1990. Physiological stress in striped bass: effect of acclimation temperature. *Aquaculture* 91:349-358.
- Davis, K.B. & Peterson, B.C. 2006. The effect of temperature, stress, and cortisol on plasma IGF-I and IGF-BPs in sunshine bass. *General and Comparative Endocrinology* 149:219-225.
- Davis, K.B., Suttle, M.A., & Parker, N.C. 1984. Biotic and abiotic influences on corticosteroid hormone rhythms in channel catfish. *Transactions of the American Fisheries Society* 113:414-421.
- de Aguiar, L.H. Moraes, G., Avilez, I.M., Altran, A.E., & Corrêa, C.F. 2004. Metabolical effects of Folidol 600 on the neotropical freshwater fish matrinxã, *Brycon cephalus*. *Environmental Research* 95:224-230.
- DeLorenzo, M.E., Scott, G.I., & Ross, P.E. 2001. Toxicity of pesticides to aquatic microorganisms: A review. *Environmental Toxicology and Chemistry* 20:84-98.
- De Mel, G.W.J.L.M.V.T.M. & Pathiratne, A. 2005. Toxicity assessment of insecticides commonly used in rice pest management to the fry of common carp, *Cyprinus carpio*, a food fish culturable in rice fields. *Journal of Applied Ichthyology* 21:146-150.
- De Silva, P.M.C.S. & Samayawardhena, L.A. 2002. Low concentrations of lorsban in water result in far reaching behavioral and histological effects in early life stages in guppy. *Ecotoxicology and Environmental Safety* 53:248-254.
- De Stasio, B.T., Jr., Hill, D.K., Kleinhans, J.M., Nibbelink, N.P., & Magnuson, J.J. 1996. Potential effects of global climate change on small north-temperate lakes: Physics, fish, and plankton. *Limnology and Oceanography* 41:1136-1149.
- Dhar, G., Barat, S., & Dhar, K.M. 2004. Influence of the organophosphate insecticide phosphamidon on lethic water. *Journal of Environmental Biology* 25:359-363.

- Dorval, J. & Hontela, A. 2003. Role of glutathione redox cycle and catalase in defense against oxidative stress induced by endosulfan in adrenocortical cells of rainbow trout (*Oncorhynchus mykiss*). *Toxicology and Applied Pharmacology* 192:191-200.
- Dorval, J., Leblond, V., Deblois, C., & Hontela, A. 2005. Oxidative stress and endocrine endpoints in white sucker (*Catostomus commersoni*) from a river impacted by agricultural chemicals. *Environmental Toxicology and Chemistry* 24:1273-1280.
- Dorval, J., Leblond, V.S., & Hontela, A. 2003. Oxidative stress and loss of cortisol secretion in adrenocortical cells of rainbow trout (*Oncorhynchus mykiss*) exposed *in vitro* to endosulfan, an organochlorine pesticide *Aquatic Toxicology* 63:229-241.
- Duchesne, J.-F. & Magnan, P. 1997. The use of climate classification parameters to investigate geographical variations in the life history traits of ectotherms, with special reference to the white sucker (*Catostomus commersoni*). *Ecoscience* 4:140-150.
- Dutta, H., Marcelino, J., & Richmonds, Ch. 1992. Brain acetylcholinesterase activity and optomotor behavior in bluegills, *Lepomis macrochirus* exposed to different concentrations of diazinon. *Archives Internationales de Physiologie, de Biochimie et de Biophysique* 100:331-334.
- Eaton, J.G. & Scheller, R.M. 1996. Effects of climate warming on fish thermal habitats in streams of the United States. *Limnology and Oceanography* 41:1109-1115.
- Eder, K.J., Leutenegger, C.M., Wilson, B.W., & Werner, I. 2004. Molecular and cellular biomarker responses to pesticide exposure in juvenile Chinook salmon (*Oncorhynchus tshawytscha*). *Marine Environmental Research* 58:809-813.
- Fairchild, W.L., Swansburg, E.O., Arsenault, J.T., & Brown, S.B. 1999. Does an association between pesticide use and subsequent declines in catch of Atlantic salmon (*Salmo salar*) represent a case of endocrine disruption? *Environmental Health Perspectives* 107:349-357.
- Fang, X., Stefan, H.G., Eaton, J.G., McCormick, J.H., & Alam, S.R. 2004a. Simulation of thermal/dissolved oxygen habitat for fishes in lakes under different climate scenarios Part 1. Cool-water fish in contiguous US. *Ecological Modelling* 172:13-37.
- Fang, X., Stefan, H.G., Eaton, J.G., McCormick, J.H., & Alam, S.R. 2004b. Simulation of thermal/dissolved oxygen habitat for fishes in lakes under different climate scenarios Part 2. Cold-water fish in contiguous US. *Ecological Modelling* 172:39-54.

- Ferrando, M.D., Sancho, E., & Andreu-Moliner, E. 1991. Comparative acute toxicities of selected pesticides to *Anguilla anguilla*. *Journal of Environmental Science and Health* 26B: 491-498.
- Ferrari, A., Anguiano, O.L., Soleño, J., Venturino, A., & Pechen de D'Angelo, A.M. 2004. Different susceptibility of two aquatic vertebrates (*Oncorhynchus mykiss* and *Bufo arenarum*) to azinphos methyl and carbaryl. *Comparative Biochemistry and Physiology* 139C:239-243.
- Flammarion, P., Migeon, B., & Garric, J. 1996. Joint effects of copper sulphate and methidathion on rainbow trout (*Oncorhynchus mykiss*) EROD and AChE activities. *Bulletin of Environmental Contamination and Toxicology* 56:440-445.
- Frank, R. & Sirons, G.J. 1979. Atrazine: Its use in corn production and its loss to stream waters in southern Ontario, 1975-1977. *Science of the Total Environment* 12:223-239.
- Frasco, M.F. & Guilhermino, L. 2002. Effects of dimethoate and beta-naphthoflavone on selected biomarkers of *Poecilia reticulata*. *Fish Physiology and Biochemistry* 26:149-156.
- Fulton, M.H. & Key, P.B. 2001. Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticide exposure and effects. *Environmental Toxicology and Chemistry* 20:37-45.
- Fulton, M.H., Moore, D.W., Wirth, E.F., Chandler, G.T., Key, P.B., Daugomah, J.W., Strozier, E.D., Devane, J., Clark, J.R., Lewis, M.A., Finley, D.B., Ellenberg, W., Karnaky, K.J., Jr., & Scott, G.I. 1999. Assessment of risk reduction strategies for the management of agricultural nonpoint source pesticide runoff in estuarine ecosystems. *Toxicology and Industrial Health* 15:201-214.
- Galbreath, P.F., Adams, N.D., & Martin, T.H. 2004. Influence of heating rate on measurement of time to thermal maximum in trout. *Aquaculture* 241:587-599.
- Geslin, M. & Auperin, B. 2004. Relationship between changes in mRNAs of the genes encoding steroidogenic acute regulatory protein and P450 cholesterol side chain cleavage in head kidney and plasma levels of cortisol in response to different kinds of acute stress in the rainbow trout (*Oncorhynchus mykiss*). *General and Comparative Endocrinology* (135) 70-80.
- Gordon, A.S., Gordon, E.P., & Dyer, B.J. 2006. Susceptibility of two fishes (*Oreochromis niloticus* and *Cyprinodon variegatus*) to *Pfiesteria shumwayae* and its associated toxin: Influence of salinity. *Harmful Algae* 5:542-547.

- Grue, C.E., Hart, A.D.M., & Mineau, P. 1991. Biological consequences of depressed brain cholinesterase activity in wildlife In: Cholinesterase-inhibiting insecticides ed. Mineau. P. Elsevier. Amsterdam.
- Habig, C., Di Giulio, R.T., & Abou-Donia, M.B. 1988. Comparative properties of channel catfish (*Ictalurus punctatus*) and blue crab (*Callinectes sapidus*) acetylcholinesterases. Comparative Biochemistry and Physiology Part C 91:293-300.
- Hamm, J.T., Wilson, B.W., & Hinton, D.E. 1998. Organophosphate-induced acetylcholinesterase inhibition and embryonic retinal cell necrosis *in vivo* in the teleost (*Oryzias latipes*). NeuroToxicology 19:853-870.
- Hasspieler, B.M., Behar, J.V., Carlson, D.B., Watson, D.E., & Di Giulio, R.T. 1994. Susceptibility of channel catfish (*Ictalurus punctatus*) and brown bullhead (*Ameiurus nebulosa*) to oxidative stress: a comparative study. Aquatic Toxicology 28:53-64.
- Hayes, W.J., Jr. 1991. Introduction. In: Handbook of pesticide toxicology Ed. Hayes, W.J., Jr. & Laws, E.R., Jr. Academic Press, Inc. London, England, 1-30.
- Hazel, J. 1969. The effect of thermal acclimation upon brain acetylcholinesterase activity of *Carassius auratus* and *Fundulus heteroclitus*. Life Sciences 8:775-784.
- Hoffmann, U. & Papendorf, T. 2006. Organophosphate poisonings with parathion and dimethoate. Intensive Care Medicine 32:464-486.
- Hogan, J.W. 1970. Water temperature as a source of variation in specific activity of brain acetylcholinesterase of bluegills. Bulletin of Environmental Contamination & Toxicology 5:347-353.
- Hontela, A. 1997. Endocrine and physiological responses of fish to xenobiotics: Role of glucocorticosteroid hormones. Reviews in Toxicology 1:1-46
- Hontela, A. 1998. Interrenal dysfunction in fish from contaminated sites: *in Vivo* and *in Vitro* assessment. Environmental Toxicology and Chemistry 17:44-48.
- Hontela, A. 2005. Adrenal toxicology: Environmental pollutants and the HPI axis. In: *Biochemistry and Molecular Biology of Fishes* (Chapter 6). Mommsen, T.P. & Moon, T.W. Eds. 332-363.
- Høy, T., Horsberg, T.E., & Wichstrøm, R. 1991. Inhibition of acetylcholinesterase in rainbow trout following dichlorvos treatment at different water oxygen levels. Aquaculture 95:33-40.

- Ihnat, J.M. & Bulkley, R.V. 1984. Influence of acclimation temperature and season on acute temperature preference of adult mountain whitefish, *Prosopium williamsoni*. *Environmental Biology of Fishes* 11:29-40.
- Intorre, L., Soldani, G., Cognetti-Varriale, A.M., Monni, G., Meucci, V., & Pretti, C. 2004. Safety of azamethiphos in eel, seabass and trout. *Pharmacological Research* 49: 171-176.
- Iwama, G.K., Thomas, P.T., Forsyth, R.B., & Vijayan, M.M. 1998. Heat shock protein expression in fish. *Reviews in Fish Biology and Fisheries* 8:35-56.
- Jarrard, H.E., Delaney, K.R., & Kennedy, C.J. 2004. Impacts of carbamate pesticides on olfactory neurophysiology and cholinesterase activity in coho salmon (*Oncorhynchus kisutch*). *Aquatic Toxicology* 69:133-148.
- Jentoft, S., Aastveit, A.H., Torjesen, P.A., & Andersen, Ø. 2005. Effects of stress on growth, cortisol, and glucose levels in non-domesticated Eurasian perch (*Perca fluviatilis*) and domesticated rainbow trout (*Oncorhynchus mykiss*). *Comparative Biochemistry and Physiology* 141A:353-358.
- Jett, D.A., Hill, E.F., Fernando, J.C., Eldefrawi, M.E., & Eldefrawi, A.T. 1993. Down-regulation of muscarinic receptors and the m3 subtype in white-footed mice by dietary exposure to parathion. *Journal of Toxicology and Environmental Health* 39:385-415.
- Johnson, J.A. & Wallace, K.B. 1987. Species-related differences in the inhibition of brain acetylcholinesterase by paraoxon and malaoxon. *Toxicology and Applied Pharmacology* 88:234-241.
- Jones, S.B., King, L.B., Sappington, L.C., Dwyer, F.J., Ellersieck, M., & Buckler, D.R. 1998. Effects of carbaryl, permethrin, 4-nonylphenol, and copper on muscarinic cholinergic receptors in brain of surrogate and listed fish species. *Comparative Biochemistry and Physiology Part C* 120:405-414.
- Kavitha, P. & Venkateswara Rao, J. 2007. Oxidative stress and locomotor behaviour response as biomarkers for assessing recovery status of mosquito fish, *Gambusia affinis* after lethal effect of an organophosphate pesticide, monocrotophos. *Pesticide Biochemistry and Physiology* 87:182-188.
- Keizer, J., D'Agostino, G., Nagel, R., Volpe, T., Gnemi, P., & Vittozzi, L. 1995. Enzymological differences of AChE and diazinon hepatic metabolism: correlation of in vitro data with selective toxicity of diazinon to fish species. *The Science of the Total Environment* 171:213-200.

- Koger, C.S., The, S.J., & Hinton, D.E. 1999. Variations of light and temperature regimes and resulting effects on reproductive parameters in Medaka (*Oryzias latipes*). *Biology of Reproduction* 61:1287-1293.
- Köprücü, S.Ş., Köprücü, K., Ural, M.Ş., İspir, Ü., & Pala, M. 2006. Acute toxicity of organophosphorus pesticide diazinon and its effects on behaviour and some hematological parameters of fingerling European catfish (*Silurus glanis* L.). *Pesticide Biochemistry and Physiology* 86:99-105.
- Kuo, C.M. & Hsieh, S.L. 2006. Comparisons of physiological and biochemical responses between milkfish (*Chanos chanos*) and grass carp (*Ctenopharyngodon idella*) to cold shock. *Aquaculture* 251:526-536.
- Lacroix, A. & Hontela, A. 2004. A comparative assessment of adrenotoxic effects of cadmium in two teleost species, rainbow trout, *Oncorhynchus mykiss*, and yellow perch, *Perca flavescens*. *Aquatic Toxicology* 67:13-21.
- Lacroix, M. & Hontela, A. 2001. Regulation of acute cortisol synthesis by cAMP-dependent protein kinase and protein kinase C in a teleost species, the rainbow trout (*Oncorhynchus mykiss*). *Journal of Endocrinology* 169:71-78.
- Lacroix, M. & Hontela, A. 2003. The organochlorine *o*'*p*-DDD disrupts the adrenal steroidogenic signalling pathway in rainbow trout (*Oncorhynchus mykiss*). *Toxicology and Applied Pharmacology* 190:197-205.
- Leblond, V.S., Bisson, M. & Hontela, A. 2001. Inhibition of cortisol secretion in dispersed head kidney cells of rainbow trout (*Oncorhynchus mykiss*) by endosulfan, an organochlorine pesticide. *General and Comparative Endocrinology* 121:48-56.
- Leblond, V.S. & Hontela, A. 1999. Effects of *in Vitro* exposures to cadmium, mercury, zinc, and 1-(2-Chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane on steroidogenesis by dispersed interrenal cells of rainbow trout (*Oncorhynchus mykiss*). *Toxicology and Applied Pharmacology* 157:16–22.
- Lermen, C.L., Lappe, R., Crestani, M., Vieira, V.P., Gioda, C.R., Schetinger, M.R.C., Baldisserotto, B., Moraes, G., & Morsch, V.M. 2004. Effect of different temperature regimes on metabolic and blood parameters of silver catfish *Rhamdia quelen*. *Aquaculture* 239:497-507.
- Levin, E.D., Swain, H.A., Donerly, S., & Linney, E. 2004. Developmental chlorpyrifos effects on hatchling zebrafish swimming behavior. *Neurotoxicology and Teratology* 26:719-723.
- Linde, A.R., Sánchez-Galán, S., Klein, D., García-Vázquez, E., & Summer, K.H. 1999. Metallothionein and heavy metals in brown trout (*Salmo trutta*) and European eel

- (*Anguilla anguilla*): A comparative study. *Ecotoxicology and Environmental Safety* 44: 168-173.
- Little, E.E. & Finger, S.E. 1990. Swimming behavior as an indicator of sublethal toxicity in fish. *Environmental Toxicology and Chemistry* 9:13-19
- Liu, J. & Pope, C.N. 1996. Effects of chlorpyrifos on high-affinity choline uptake and [³H]Hemicholinium-3 binding in rat brain. *Fundamental and Applied Toxicology* 34:84-90.
- Macek, K.J., Hutchinson, C., & Cope, O.B. 1969. The effect of temperature on the susceptibility of bluegills and rainbow trout to selected pesticides. *Bulletin of Environmental Contamination & Toxicology* 4:174-183.
- Marrs, T.C. 1993. Organophosphate poisoning. *Pharmacology and Therapeutics* 58:51-66.
- McHugh, P., & Budy, P. 2005. An experimental evaluation of competitive and thermal effects of brown trout (*Salmo trutta*) and Bonneville cutthroat trout (*Oncorhynchus clarkia utah*) performance along an altitudinal gradient. *Canadian Journal of Fisheries and Aquatic Sciences* 62:2784-2795.
- Milligan, C.L. 2003. A regulatory role for cortisol in muscle glycogen metabolism in rainbow trout *Oncorhynchus mykiss* Walbaum. *Journal of Experimental Biology* 206:3167-3173.
- Mommsen, T.P., Vijayan, M.M., & Moon, T.W. 1999. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. *Reviews in Fish Biology and Fisheries* 9:211-268.
- Morgan, I.J., D'Cruz, L.M., Dockray, J.J., Linton, T.K., McDonald, D.G., & Wood, C.M. 1998. The effects of elevated winter temperature and sub-lethal pollutants (low pH, elevated ammonia) on protein turnover in gill and liver of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiology and Biochemistry* 19:377-389.
- Morrison, J., Quick, M.C., & Foreman, M.G.G. 2002. Climate changes in the Fraser River watershed: flow and temperature projections. *Journal of Hydrology* 263:230-244.
- Myrick, C.A. & Cech, J.J., Jr. 2004. Temperature effects on juvenile salmonids in California's central valley: what we don't know? *Reviews in Fish Biology and Fisheries* 14:113-123.
- Norris, D.O. 1997. The mammalian adrenal glands and Comparative aspects of vertebrate adrenals: Cortical and chromaffin cells. In: *Vertebrate Endocrinology*, edited by D.O. Norris, New York, Academic Press, pp. 299-355.

- O'Brien, R.D. 1976. Acetylcholinesterase and its inhibition in: Insecticide biochemistry and physiology ed. Wilkinson, C.F. Plenum Press New York 271-293.
- Patiño, R., Bradford, C.S., & Shreck, C.B. 1986. Adenylate cyclase activators and inhibitors, cyclic nucleotide analogs, and phosphatidylinositol: effects on interrenal function of coho salmon (*Oncorhynchus kisutch*) *in Vitro*. General and Comparative Endocrinology 63:230-235.
- Pavlov, D.D., Chuiko, G.M. Gerassimov, Y.V. & Tonkopyi, V.D. 1992. Feeding behavior and brain acetylcholinesterase activity in bream (*Abramis brama* L.) as affected by DDVP, an organophosphorus insecticide. Comparative Biochemistry and Physiology 103C:563-568.
- Payne, J.F., Mathieu, A., Melvin, W., & Fancey, L.L. 1996. Acetylcholinesterase, an old biomarker with a new future? Field trials in association with two urban rivers and a paper mill in Newfoundland. Marine Pollution Bulletin 32:225-231.
- Pehkonen, S.O. & Zhang, Q. 2002. The degradation of organophosphorus pesticides in natural waters: A critical review. Critical Reviews in Environmental Science and Technology 32:17-72.
- Phillips, T.A., Summerfelt, R.C., & Atchison, G.J. 2002. Environmental, biological, and methodological factors affecting cholinesterase activity in walleye (*Stizostedion vitreum*). Archives of Environmental Contamination and Toxicology 43:75-80.
- Pope, C.N. 1999. Organophosphorus pesticides: Do they all have the same mechanism of toxicity? Journal of Toxicology and Environmental Health 2B:161-181.
- Rajagopal, P.K. 1979. The embryonic development and the thermal effects on the development of the mountain whitefish, *Prosopium williamsoni* (Girard). Journal of Fish Biology 15:153-158.
- Rahel, F.J., Keleher, C.J., & Anderson, J.L. 1996. Potential habitat loss and population fragmentation for cold water fish in the North Platte River drainage of the Rocky Mountains: Response to climate warming. Limnology and Oceanography 41:1116-1123.
- Ratageri, R.H., Taranath, T.C., & Lakshman, H.C. 2006. Toxicity of dimethoate on primary productivity of a lentic aquatic ecosystem: A microcosm approach. Bulletin of Environmental Contamination and Toxicology 76:373-380.
- Raupach, M.R. Briggs, P.R. Ahmad, N. & Edge, V.E. 2001a. Endosulfan transport: II. Modeling airborne dispersal and deposition by spray and vapor. Journal of Environmental Quality 30:729-740.

- Raupach, M.R., Briggs, P.R., Ford, P.W., Leys, J.F., Muschal, M. Cooper, B. & Edge, V.E. 2001b. Endosulfan transport: I. Integrative assessment of airborne and waterborne pathways. *Journal of Environmental Quality* 30:714-728.
- Saglio, P., Bretaud, S., Rivot, E., & Olsén, K.H. 2003. Chemobehavioral changes induced by short-term exposures to prochloraz, nicosulfuron, and carbofuran in goldfish. *Archives of Environmental Contamination and Toxicology* 45:515-524.
- Samuel, M. & Sastry, K.V. 1989. *In vivo* effect of monocrotophos on the carbohydrate metabolism of the freshwater snake head fish, *Channa punctatus*. *Pesticide Biochemistry and Physiology* 34:1-8.
- Sancho, E., Ferrando, M.D., Andreu, E. 1997. Sublethal effects of an organophosphate insecticide on the European eel, *Anguilla anguilla*. *Ecotoxicology and Environmental Safety* 36:57-65.
- Sancho, E., Ferrando, M.D., Fernández, C., & Andreu, E. 1998. Liver energy metabolism of *Anguilla anguilla* after exposure to fenitrothion. *Ecotoxicology and Environmental Safety* 41:168-175.
- Sandahl, J.F., Baldwin, D.H., Jenkins, J.J., & Scholz, N.L. 2005. Comparative thresholds for acetylcholinesterase inhibition and behavioral impairment in coho salmon exposed to chlorpyrifos. *Environmental Toxicology and Chemistry* 24:136-145.
- Sandahl, J.F. & Jenkins, J.J. 2002. Pacific steelhead (*Oncorhynchus mykiss*) exposed to chlorpyrifos: Benchmark concentration estimates for acetylcholinesterase inhibition. *Environmental Toxicology and Chemistry* 21:2452-2458.
- Sapozhnikova, Y., Bawardi, O., & Schlenk, D. 2004. Pesticides and PCPs in sediments from fish from the Salton Sea, California, USA. *Chemosphere* 55:797-809.
- Scholz, N.L., Truelove, N.K., French, B.L., Berejikian, B.A., Quinn, T.P., Casillas, E., & Collier, T.K. 2000. Diazinon disrupts antipredator and homing behaviors in Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* 57:1911-1918.
- Scholz, N.L., Truelove, N.K., Labenia, J.S., Baldwin, D.H., & Collier, T.K. 2006. Dose-additive inhibition of Chinook salmon acetylcholinesterase activity by mixtures of organophosphate and carbamate insecticides. *Environmental Toxicology and Chemistry* 25:1200-1207.
- Schreck, C.B., Contreras-Sanchez, W., & Fitzpatrick, M.S. 2001. Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture* 197:3-24.
- Selvi, M., Sarikaya, R., Erkoç, F. & Koçak, O. 2005. Investigation of acute toxicity of chlorpyrifos-methyl on guppy *Poecilia reiculata*. *Chemosphere* 60:93-96.

- Shankland, D.L. 1976. The nervous system: Comparative physiology and pharmacology in: Insecticide biochemistry and physiology ed. Wilkinson, C.F. Plenum Press New York 229-270.
- Shao-nan, L. & De-fang, F. 1996. Correlation between biochemical parameters and susceptibility of freshwater fish to malathion. *Journal of Toxicology and Environmental Health* 48:413-418.
- Shaonan, L., Xianchuan, X., Guonian, Z., & Yajun, T. 2004. Kinetic characters and resistance to inhibition of crude and purified brain acetylcholinesterase of three freshwater fishes by organophosphates. *Aquatic Toxicology* 68:293-299.
- Singh, S.K., Tripathi, P.K., Yadav, R.P., Singh, D., & Singh, A. 2004. Toxicity of malathion and carbaryl pesticides: Effects on some biochemical profiles of the freshwater fish *Colisa fasciatus*. *Bulletin of Contamination and Toxicology* 72:592-599.
- Smith, D.L. & Almaraz, J.J. 2004. Climate change and crop production: contributions, impacts, and adaptations. *Canadian Journal of Plant Pathology* 26:253-266.
- Staurnes, M. 2001. Difference between Atlantic halibut (*Hippoglossus hippoglossus* L.) and turbot (*Scophthalmus maximus* L.) in tolerance to acute low temperature exposure. *Aquaculture Research* 32:251-255.
- Stefan, H.G., Hondzo, M., & Fang, X. 1993. Lake water quality modeling for projected future climate scenarios. *Journal of Environmental Quality* 22:417-431.
- Stenersen, J. 2004. Chemical pesticides: Mode of action and toxicology. CRC Press. Boca Ration, Florida.
- Stocco, D.M. 2000. Intermitochondrial cholesterol transfer. *Biochimica et Biophysica Acta* 1486:184-197.
- Stocco, D.M., Wang, X., Jo, Y., & Manna, P.R. 2005. Multiple signaling pathways regulating steroidogenesis and steroidogenic acute regulatory protein expression: more complicated than we thought. *Molecular Endocrinology* 19:2647-2659.
- Strange, R.J. 1980. Acclimation temperature influences cortisol and glucose concentrations in stressed channel catfish. *Transactions of the American Fisheries Society* 109:289-303.
- Strange, R.J., Schreck, C.B., & Golden, J.T. 1977. Corticoid stress response to handling and temperature in salmonids. *Transaction of the American Fisheries Society* 106:213-218.

- Strum, A., Wogram, J., Hansen, P.-D., & Liess, M. 1999. Potential use of cholinesterase in monitoring low levels of organophosphates in small streams: Natural variability in three-spined stickleback (*Gasterosteus aculeatus*) and relation to pollution. *Environmental Toxicology and Chemistry* 18:194-200.
- Sweilum, M.A. 2006. Effect of sublethal toxicity of some pesticides on growth parameters, haematological properties and total production of Nile tilapia (*Oreochromis niloticus* L.) and water quality of ponds. *Aquaculture Research* 37:1079-1089.
- Szabó, A., Nemcsók, J., Kása, P., & Budai, D. 1991. Comparative study of Acetylcholine synthesis in organs of freshwater teleosts. *Fish Physiology and Biochemistry* 9:93-99.
- Tate, K.W. & Lancaster, D.L. 2007. Assessment of thermal stratification within stream pools as a mechanism to provide refugia for native trout in hot, arid rangelands. *Environmental Monitoring and Assessment* 124:289-300.
- Thangavel, P., Sumathiral, K., Karthikeyan, S., & Ramaswamy, M. 2005. Endocrine response of the freshwater teleost, *Sarotherodon mossambicus* (Peters) to dimecron exposure. *Chemosphere* 61:1083-1092.
- Tilak, K.S., Veeraiah, K., & Koteswara Rao, D. 2005. Biochemical changes induced by chlorpyrifos, an organophosphate compound in sublethal concentrations to the freshwater fish *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*. *Journal of Environmental Biology* 26:341-347.
- Trim, A.H. & Marcus, J.M. 1990. Integration of long-term fish kill data with ambient water quality monitoring data and application to water quality management. *Environmental Management* 14:389-396.
- Tuduri, L. Harner, T., Blanchard, P., Li, Y.-F., Poissant, L., Waite, D.T., Murphy, C., & Belzer, W. 2006a. A review of currently used pesticides (CUPs) in Canadian air and precipitation : Part 1: Lindane and endosulfans. *Atmospheric Environment* 40: 1563-1578.
- Tuduri, L. Harner, T., Blanchard, P., Li, Y.-F., Poissant, L., Waite, D.T., Murphy, C., & Belzer, W. 2006b. A review of currently used pesticides (CUPs) in Canadian air and precipitation : Part 2: Regional information and perspectives. *Atmospheric Environment* 40: 1563-1578.
- van Dijk, P.L.M., Tesch, C., Hardewig, I., & Pörtner, H.O. 1999. Physiological disturbances at critically high temperatures: a comparison between stenothermal antarctic and eurythermal temperate eelpouts (*Zoarctidae*). *The Journal of Experimental Biology* 202:3611-3621.

- Van Dolah, R.F., Maier, P.P., Fulton, M.H., & Scott, G.I. 1997. Comparison of azinphosmethyl toxicology to juvenile red drum (*Sciaenops ocellatus*) and the mummichog (*Fundulus heteroclitus*). *Environmental Toxicology and Chemistry* 16:1488-1493.
- Viant, M.R., Werner, I., Rosenblum, E.S., Gantner, A.S., Tjeerdema, R.S., & Johnson, M.L. 2003. Correlation between heat-shock protein induction and reduced metabolic condition in juvenile steelhead trout (*Oncorhynchus mykiss*) chronically exposed to elevated temperature. *Fish Physiology and Biochemistry* 29:159-171.
- Vittozzi, L. & De Angelis, G. 1991. A critical review of comparative acute toxicity data on freshwater fish. *Aquatic Toxicology* 19:167-204.
- Waite, D.T., Grover, R., Westcott, N.D., & Irvine, D.G. 1995. Atmospheric deposition of pesticides in a small southern Saskatchewan watershed. *Environmental Toxicology and Chemistry* 14:1171-1175.
- Wang, J., Grisle, S., & Schlenk, D. 2001. Effects of salinity on aldicarb toxicity in juvenile rainbow trout (*Oncorhynchus mykiss*) and striped bass (*Morone saxatilis* X *chrysops*). *Toxicological Sciences* 64:200-207.
- Waring, C.P. & Moore, A. 2004. The effects of atrazine on Atlantic salmon (*Salmo salar*) smolts in fresh water and sea water transfer. *Aquatic Toxicology* 66:93-104.
- Weber, T.E. & Bosworth, B.G. 2005. Effects of 28 day exposure to cold temperature or feed restriction on growth, body composition, and expression of genes related to muscle growth and metabolism in channel catfish. *Aquaculture* 246:483-492.
- Weiss, C.M. 1958. The determination of cholinesterase in the brain tissue of three species of fresh water fish and its inactivation in Vivo. *Ecology* 39:194-199.
- Wheelock, C.E., Eder, K.J., Werner, I., Huang, H., Jones, P.D., Brammell, B.F., Elskus, A.A., & Hammock, B.D. 2005. Individual variability in esterase activity and CYP1A levels in Chinook salmon (*Oncorhynchus mykiss*) exposed to esfenvalerate and chlorpyrifos. *Aquatic Toxicology* 74:172-192.
- Whitehead, A., Anderson, S.L., Ramirez, A., & Wilson, B.W. 2005. Cholinesterase in aquatic biomonitoring: Assay optimization and species-specific characterization for a California native fish. *Ecotoxicology* 14:597-606.
- Zinkl, J.G., Shea, P.J., Nakamoto, R.J., & Callman, J. 1987. Technical and biological considerations for the analysis of brain cholinesterase of rainbow trout. *Transactions of the American Fisheries Society* 116:570-573.

Chapter 2.

The physiological stress response indicators in whitefish (*Prosopium williamsoni*) and suckers (*Catostomus sp.*) sampled along a gradient of temperature and agricultural chemicals in the Oldman river, Alberta.

2.1 Introduction

Rivers are complex systems where habitats of different species overlap. Species that occupy the same niche may share resources and co-exist, or they may compete with each other. A species better suited for a particular habitat can cause a decline in another species' population. Anthropogenic factors such as physical structures (dams and weirs), water withdrawals or loading of toxicants can cause changes in the aquatic environment (Cormier et al. 2000), potentially stressing and adversely affecting the organisms that live there. Differences in sensitivity to stressors between species can lead to the extirpation of some species and the invasion of others. Species-specific responses of fish have been documented with respect to toxicity (Gordon et al. 2006), acetylcholinesterase (AChE) inhibition (Davies et al. 1994, Fulton & Key 2001), metal distribution and metallothionein concentration (Linde et al. 1999), stress and the stress response (Lacroix & Hontela 2004, Jentoft et al. 2005), and vulnerability to oxidative stress (Hasspieler et al. 1994). Differences in tolerance to heat (Van Dijk et al. 1999) and cold shock (Kuo & Hsieh 2006) have been also reported; fish are classified as cool- or cold-water species according to their thermal preference. Cool- and cold-water fish species often share the same environment, particularly when their movements are restricted by dams and weirs. As climate changes and water temperature increases, temperature sensitive species may be threatened. Cold-water fish species in rivers draining agricultural areas in temperate

zones are particularly vulnerable to pesticides because they are also exposed to high temperatures.

In the agricultural drainages, fish are exposed to short-term spikes of pesticides during spraying and rain events, as well as to chronic exposure in areas where streams and canals drain sprayed areas (Frank & Sirons 1979, Morrison & Wells 1981). Exposure to organophosphate (OP) and carbamate (CB) pesticides lower the activity of the enzyme AChE. Low levels of AChE in fish cause changes in behaviour (Beauvais et al. 2000, Brewer et al. 2001, de Aguiar et al. 2004, Sandahl et al. 2005), metabolism (Samuel & Sastry 1989, de Aguiar et al. 2004), feeding (Pavlov et al. 1992) and reproduction (Bhattacharya 1993). Agricultural chemicals, including pesticides have also been shown to influence the physiological stress response and cortisol secretion in fish (Dorval et al. 2005, Teles et al. 2007; Thangravel et al. 2005). Natural stressors such as increases in temperature can modify the toxicity of pesticides; warmer water temperature increased the toxicity of endosulfan (Capkin et al. 2006) and methicarb (Altinok et al. 2006) to rainbow trout. Moreover, an increase in global temperature may lead to invasions of new and more abundant agricultural pests, thus increasing the amount of pesticides being used (Moore & Folt 1993, Bloomfield et al. 2006).

Fish have precise thermal requirements and will select habitats that are near their optimal temperature (Neill & Magnuson 1974, Dill 1987); if necessary, they will expand their range of temperature in response to food availability (Brett 1971), predators, competition, or social dominance (Neill & Magnuson 1974, Beitinger & Magnuson 1975, Dill 1987). Increasing water temperature is considered a widespread and problematic stressor in salmonids on the West Coast (Sullivan et al. 2000); displacement of cold-

water, northern, high latitude and high altitude species of fish is predicted in many global warming models (De Stasio et al. 1996, Daufresne et al. 2003). Water temperatures outside the thermal preference range, specifically warmer water temperatures, can cause thermal stress which depletes energy reserves (Brett 1971, Viant et al. 2003), decreases growth rates (Brett 1971, Bartell et al. 1986), impairs reproduction (Daufresne et al. 2003) and changes behaviour (Sullivan et al. 2000).

Fish in anthropogenically impacted regions may be exposed to many different concurrent stressors, including loading of agricultural chemicals such as pesticides and fertilizers, increasing water temperatures, and habitat alterations (dams, weirs, and water withdrawals). The physiological stress response is activated when homeostasis is challenged and there are changes to tissue and organ function in an attempt to maintain internal stability (Wendelaar Bonga 1997, Schreck et al. 2001). Physiological indicators of stress such as plasma cortisol and glucose levels, and liver glycogen are often used to determine if fish are stressed and if the hypothalamo-pituitary-interrenal axis functions normally (Hontela 1998). The type and duration of a stressor can have an impact on the stress response: acute exposures to a wide range of chemical pollutants increase plasma cortisol in fish (Hontela 1997) while chronic exposure may disrupt cortisol synthesis. Agricultural chemicals (Dorval et al. 2005) and metals such as Cd, Cu, and Zn (Laflamme et al. 2000, Lacroix & Hontela 2004, Lappivaara & Marttien 2005), can impair the stress response resulting in low levels of cortisol. Pesticides and increasing water temperatures are stressors potentially affecting fish in many regions in North America.

Water temperatures in the Experimental Lake Area of Northern Ontario, Canada have risen by 2°C in the last 20 years (Schindler et al. 1996). Global water temperatures are projected to rise by 1-7°C (depending on type and location of water body) when a doubling of atmospheric CO₂ occurs (Stefan et al. 1993, De Stasio et al. 1996, Eaton & Scheller, 1996). A combination of global climate change and continued runoff of agricultural chemicals may have disastrous effects on fisheries and may decrease the habitat for cold-water and sensitive fish. Yet our understanding of the interactions between temperature and chemical stressors on fish physiological fitness is limited.

The objective of this study is to determine how multiple stressors (pesticides and warm temperature) affect the physiological status and the stress response in cold-water and cool-water fish. Plasma cortisol and glucose, liver glycogen, and condition factor were measured as the physiological stress response endpoints, and plasma AChE activity was used as an indicator of exposure to OP and CB pesticides. It is hypothesised that species-specific differences to these stressors exist and that at temperature-impacted sites, the cold-water fish species will exhibit an increased stress response compared to cool-water fish.

2.2 Materials and Methods

2.2.1 Description of the sites and fish species

Fish were sampled at six sites along the Oldman River (Figure 2.1), in southern Alberta, Canada during spring and/or summer 2004 and 2005 (Table 2.1). The Oldman River originates in the Rocky Mountains and flows east, draining an area of intensive grain and livestock production. Runoff from these operations contains agricultural



Figure 2.1. Location of sampling sites in the Oldman River Basin, Southern Alberta. Squares represent site locations, the x is the location of the Oldman Dam and arrows indicate flow direction, with Summerview the most upstream site and Pavan the most downstream site. Figure modified from Oldman Watershed Council (2006)

Table 2.1. Characteristics of sites sampled in the Oldman River, Alberta.

| Site | Distance from Oldman dam (km) | Latitude and longitude | Date | Depth* (m) | Velocity* (m/s) | Flow* (m ³ /s) |
|-------------|-------------------------------|------------------------|-------------|------------|-----------------|---------------------------|
| Summerview | 7 | 49°33'N, 113°49'W | Summer 2005 | 0.6 | 0.4 | 3.6-6 |
| | | | Spring 2005 | 0.4 | 0.3 | 2.0 |
| Fort McLeod | 70 | 49°44'N, 113°13'W | Summer 2005 | 0.3-0.7 | 0.1-0.6 | 0.9-5.2 |
| | | | Spring 2005 | 0.4-0.7 | 0-0.5 | |
| Pearce Road | 105 | 49°48'N, 113°13'W | Summer 2005 | 0.4-0.5 | 0.1-0.7 | 0.5-5.5 |
| | | | Spring 2005 | 0.3-1.1 | 0.1-0.5 | 0.5-5.0 |
| Popson Park | 160 | 49°44'N, 112°51'W | Summer 2005 | 0.7 | | |
| | | | Spring 2005 | 0.4-0.6 | 0.1-0.4 | 0.6-3.0 |
| Pavan Park | 175 | 49°38'N, 112°51'W | Summer 2005 | 0.4-0.8 | 1.4-1.6 | 9.6-31.5 |

*Measurements taken at points in the river where fish were seined

chemicals including pesticides that are detected in the water (Table 2.2). Moreover, the river exhibits a west to east temperature gradient, with higher temperatures in the lower reaches (Figure 2.2). Whitefish (*Prosopium williamsoni*) and suckers (*Catostomus* sp.) were sampled for this study because both species are found within the Oldman River and they have different species-specific temperature optima. Whitefish represent a cold-water fish species with a maximum temperature tolerance of 23.1°C (Eaton & Scheller 1996) and a prespawning thermal preference range between 12.8°C to 17.7°C [upper temperature preference (UTP)] (Ihnat & Bulkley 1984). Suckers represent a cool-water species with a maximum temperature tolerance of 27.4°C (Eaton & Scheller 1996) and a thermal preference range of 18.3°C to 24°C (UTP) (Eaton et al. 1995). The UPTs for sucker and whitefish are indicated in Figure 2.2.

2.2.2 Capture and sampling of the fish

Fish were captured by a purse seine (30 x 1.8 m, 0.4 cm mesh) between 10:00 h and 13:30 h each day, at sampling periods indicated in Table 2.1. Fish were transferred from the seine to a floating enclosure (0.68 x 1 x 0.47 m) and were kept in the enclosure until 14:00 h when they were removed in groups of 3 to 5. Fish were anaesthetized in a solution of 0.15 g/L of tricaine methanesulphonate (MS-222) and a blood sample was taken from the caudal vasculature with a 1 or 3 mL heparized syringe. Plasma was separated at 13000 rpm for 5 minutes and frozen in liquid nitrogen. The fish were euthanized by spinal transection, weights and lengths recorded, and the liver was dissected and frozen in liquid nitrogen. Plasma and liver samples were kept frozen at -80°C until analysis.

Table 2.2. Concentrations of chemicals in water detected at sites situated along the Oldman River, Alberta.

| Compound | | Summerview | Fort MacLeod | Monarch | Lethbridge | Popson Park | Hwy 36 |
|-------------------------------|----------------------|-------------------------|--------------|-----------|------------------|-------------|-----------------|
| | 2,4-D | <0.005(04) ^a | | | 0.005-0.026(04) | | 0.006-0.082(04) |
| Herbicide (µg/L) | Dichlorprop (2,4-DP) | <0.005(04) | | | <0.005-0.007(04) | | <0.005(04) |
| | MCPA | <0.005(04) | | | <0.005-0.01(04)* | | 0.003-0.022(04) |
| | MCPP (Mecoprop) | <0.005(04) | | | <0.005(04) | | 0.004-0.018(04) |
| | Dicamba (Banvel) | <0.005(04) | | | <0.005(04) | | <0.005-0.006 |
| Insecticide (µg/L) | Dimethoate (Cygon) | <0.05(04) | <0.05(98) | <0.05(98) | <0.05(04)* | <0.05(99) | <0.05(04)* |
| | Cholesterol | 74.70(05) | | | | 108.15(05) | 217.98(05) |
| | Desmosterol | 28.46(05) | | | | 36.34(05) | 51.58(05) |
| | Cholestanol | 3.61(05) | | | | 12.26(05) | 36.27(05) |
| Other Chemicals† (ng/L) | 7-Ketocholesterol | 1.65(05) | | | | 7.27(05) | 9.90(05) |
| | β-Sitosterol | 40.56(05) | | | | 48.62(05) | 118.74(05) |
| | Campesterol | 9.43(05) | | | | 10.12(05) | 16.75(05) |
| | Stigmastanol | 2.47(05) | | | | 10.64(05) | 33.30(05) |
| | Stigmasterol | 19.80(05) | | | | 42.01(05) | 118.20(05) |
| | Fucosterol | 65.18(05) | | | | 111.63(05) | 208.14(05) |

^a Year of sampling indicated in bracket; 2004 as (04) and 2005 as (05)

† Samples from Jeffries 2007

Organophosphates and Carbamates measured by Alberta Environment were below the detection limit for that pesticide at all sites : Chlorpyrifos-ethyl, Diazinon and Phorate <0.005µg/L, Parathion <0.01 µg/L, Terbufos <0.03 µg/L, Malathion and Triallate <0.05 ug/L, Ethion <0.1 µg/L, Disulfoton and Gluthion <0.2 µg/L




Figure 2.2. Mean water temperatures for each site during sampling periods indicated by black bars below the dates in A) Early Summer and Late Summer 2004 B) Spring and Late summer 2005. Data for Summerview, Lethbridge and Highway 36 received from Alberta Environment. Upper thermal preference (UTP) for sucker and for whitefish are indicated by the lines on the graph for sucker (-.-.-) and whitefish (- - -).

2.2.3 Acetylcholinesterase (AChE)

The AChE activity was measured with an assay modified from Ellman et al. (1961) and Chuiko (2000). Plasma (2 μ L) was pipetted into a microplate and 120 μ L of ISO-OMPA (Sigma Tetraisopropyl pyrophosphoramidate, T1505) was added. The samples were incubated at room temperature for 10 minutes and 10 μ L of DTNB (Sigma 5,5'-Dithiobis [2-nitro-benzoic acid], D8130) and 10 μ L of Acetylthiochlorine Iodine (Sigma minimum 98% TLC, A5751) were added, followed by a 10 minute incubation at room temperature. The microplate was read at 405 nm in a microplate reader every 2 minutes for 10 minutes. Concentration of AChE for each sample was measured from the slope; internal standards (Normal Serum Control, TC-TROL[N], Teco Diagnostics) were used to ensure the accuracy of the assay was maintained.

2.2.4 Physiological Stress Response

Concentrations of cortisol in the plasma were determined with diagnostic kits (MP Biomedicals Diagnostics Division 07-221102). Assay characteristics and accuracy were verified, as described previously (Hontela et al. 1995). Plasma glucose was determined with a colorimetric assay using the GOD-PAP reagent (Roche 1929526) at 512nm. Liver glycogen was determined using a method modified from Bleau et al. (1996). Liver samples were digested in KOH (1N) at room temperature; the sample was then precipitated with 1.5N acetic acid and centrifuged at 13000 RPM for 10 minutes. Glycogen from the livers and standards (rabbit glycogen, Sigma, G8876-1G) was hydrolyzed to glucose with aminoglycosidase (Sigma A-7255) in a water bath at 37°C for 30 minutes. Glucose content was measured using the colorimetric assay described above.

2.2.5 Statistical Analysis


The statistical program JMP IN version 5.1 was used to perform statistical tests at $p < 0.05$. One way ANOVA was used for single factor comparisons, a t-test was used to compare species at each site. Significant differences between means were determined using Tukey-Kramer HSD post hoc test and log-normal data were normalized using a log transformation.

2.3 Results

2.3.1 Pesticide Exposure

Water concentrations of herbicides and other chemicals in the Oldman river increased along a west to east transect in 2004 and 2005 (Table 2.2), however several of the OP and CB pesticides that were measured were below the detection limits (0.005-0.1 $\mu\text{g}/\text{l}$ depending on the pesticide) (Table 2.2). Plasma AChE activities, used as indicators of exposure to OP and CB pesticides, are presented in Fig. 2.3. At all sites and all sampling times, whitefish had significantly lower plasma AChE activity ($p < 0.05$, t-test) compared to suckers (Figure 2.3A & 2.3B). In spring 2005, AChE activities in whitefish were lower at downstream sites while plasma AChE in suckers were not significantly different among sites ($p > 0.05$, ANOVA) (Figure 2.3A). Plasma AChE activity in late summer 2005 was higher at downstream sites in both suckers and whitefish (Figure 2.3B).

A) Spring 2005



B) Late Summer 2005




Figure 2.3. Plasma AChE (mean±SE) in whitefish and suckers sampled at sites along the Oldman River in A) spring 2005 B) late summer 2005. Lines under sites represent sites that are within the thermal preference of whitefish. Capital letters represent significant differences among sites for suckers and lower case letters represent significant differences among sites for whitefish (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Asterisk represent significant differences between species at a site (Student's t-test $\alpha=0.05$). Sites with only one bar indicate that only one species was caught at that site.


2.3.2 Temperature Regimes

Water temperatures fluctuated throughout the sampling period and were always below the maximum temperature tolerance for whitefish and suckers, 23.1°C and 27.4°C respectively (Eaton & Scheller 1996). However water temperatures did exceed the UTP for whitefish (17.7°C) during: early summer 2004 (Figure 2.2A & 2.2B) at Pearce Rd, Monarch (2nd day sampling), Popson, and Pavan (data extrapolated from Alberta Environment), late summer 2004 at Pearce Rd, Monarch, Popson, and Pavan, and in late summer 2005 at Pearce Rd, Popson, and Pavan. In early summer 2004, the water temperature was abnormally high during sampling at Pearce Rd, ranging from 23.0 to 24.8°C. In spring 2005 all sites where whitefish were sampled were within their thermal preference (Figure 2.2B). Whitefish were not found at Popson in 2004, late summer 2005, or at Pavan in any of the sampling periods. Suckers were not present at Summerview in any of the sampling periods.


2.3.4 Physiological Stress Response

Plasma cortisol levels are presented in Figure 2.4. Plasma cortisol levels in whitefish were higher compared to suckers at 7 sampling times (significantly higher at $p < 0.05$ at 6 sampling times) out of a total of 11 sampling times. Suckers had significantly higher ($p < 0.05$) plasma cortisol compared to whitefish 3 out of 11 sampling times and always at Ft MacLeod. In 2004, there were no significant differences in plasma cortisol for whitefish sampled at different sites but in early summer 2004 whitefish at Pearce Rd have significantly higher plasma cortisol than at all other sites (Figure 2.4A). There was no significant difference ($p > 0.05$) in plasma cortisol in suckers in 2004 (Figure


A) Early Summer 2004



B) Late Summer 2004



C) Spring 2005



D) Late Summer 2005




Figure 2.4. Plasma cortisol (mean±SE) in whitefish and suckers sampled at sites along the Oldman River in A) early summer 2004, B) late summer 2004, C) spring 2005, and D) late summer 2005. Numbers in the bars indicate the sample size for each site. Lines under sites represent sites within the thermal preference of whitefish. Capital letters represent significant differences among sites for suckers and lower case letters represent significant differences among sites for whitefish (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Asterisks represent significant differences between species at a site (t-test $\alpha=0.05$). Sites with only one bar indicate that only one species was present at that site.

2.4B) between sites. In 2005, plasma cortisol of whitefish at downstream sites was higher than upstream (Figure 2.4C & 2.4D); plasma cortisol in suckers also increased at downstream sites in late summer 2005 (Figure 2.4D).

The concentrations of liver glycogen (Fig. 2.5), measured as an indicator of energy reserves, were significantly lower ($p < 0.05$, t-test) in whitefish compared to suckers at all sites and all sampling times, except at Ft. MacLeod in spring 2005 where the means were not different (Fig. 2.5C). In both fish species sampled in late summer 2004 (Fig. 2.5B), spring 2005 (Fig. 2.5C), and late summer 2005 (Fig. 2.5D), liver glycogen reserves decreased along a west to east gradient. This trend was however not observed in early summer 2004 (Fig. 2.5A).

Plasma glucose levels of whitefish and suckers did not show any site- or species-related trends (Table 2.3). In whitefish there was a decrease in plasma glucose levels at downstream sites in 2005 but suckers did not demonstrate this trend at any sampling period. Condition factor for either whitefish or suckers did not show any observable patterns for any sample period (Table 2.4).

2.4 Discussion

2.4.1 Exposure

The Oldman River flows east from the Rocky Mountains until it joins the Bow River to form the South Saskatchewan River just east of Taber, AB. The temperature of the Oldman River rises as it flows toward the South Saskatchewan River, creating an east to west temperature gradient. The distribution of fish in the Oldman River reflects this temperature gradient, as mainly cold-water species, such as rainbow trout (*Oncorhynchus*




Figure 2.5. Liver glycogen (mean±SE) in whitefish and suckers sampled at sites along the Oldman River in A) early summer 2004, B) late summer 2004, C) spring 2005, and D) late summer 2005. Numbers in the bars indicate the sample size for each site. Lines under sites represent sites within the thermal preference of whitefish. Capital letters represent significant differences among sites for suckers and lower case letters represent significant differences among sites for whitefish (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Asterisk represent significant differences between species at a site (t-test $\alpha=0.05$). Sites with only one bar indicate that only one species was present at that site.

Table 2.3. Plasma Glucose, mean±SE(n), in whitefish and suckers sampled at sites along the Oldman River in early summer 2004, late summer 2004, spring 2005, late summer 2005.

| Sampling Period | Species | Glucose (mg/mL) | | | | | |
|-----------------|-----------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------------|
| | | Sum | Ft.M | Per | Mon | Pop | Pav |
| E. Summer '04 | Whitefish | 1.84±0.05 (33) a | 1.85±0.06(27) a* | 1.55±0.10(34) b | 1.93±0.16(9) ab* | - | - |
| | Sucker | - | 1.57±0.11(31) AB | 1.64±0.12(33) A | 1.40±0.09(23) AB | 1.13±0.10(21) B | 2.07±0.19(18) A |
| L. Summer '04 | Whitefish | 1.63±0.07(24) ab | 1.42±0.06(25) b | 1.76±0.10(15) a* | 1.56±0.08(15) ab | - | - |
| | Sucker | - | 1.98±0.12(22) A* | 1.11±0.13(8) BC | 1.34±0.14(19) B | 1.00±0.06(57) C | 1.32±0.07(35) B |
| Spring '05 | Whitefish | | 1.55±0.08(10) a | 1.51±0.09(19) a | | 1.15±0.11(12) b | - |
| | Sucker | | 1.44±0.08(25) A | 1.39±0.13(22) A | | 1.55±0.11(20) A* | 1.86±0.26(9) A |
| L. Summer '05 | Whitefish | 2.06±0.09(14) a | 1.68±0.07(18) b | 1.63±0.14(15) b | | - | - |
| | Sucker | - | 1.92±0.16(20) A | 1.88±0.11(20) A | | 1.93±0.10(18) A | 2.10±0.14(16) A |

Lowercase letters represent significant differences within a sampling period for whitefish and capital letters represent significant differences within a sampling period for suckers (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Asterisk represents a significant difference between suckers and whitefish at a site (T-test $\alpha=0.05$)

- species not caught during sampling period

Table 2.4. Condition factor, mean±SE(n), in whitefish and suckers sampled at sites along the Oldman River in early summer 2004, late summer 2004, spring 2005, late summer 2005.

| Sampling periods | Species | Condition Factor | | | | | |
|------------------|-----------|---------------------|---------------------|--------------------|---------------------|--------------------|---------------------|
| | | Sum | Ft.M | Per | Mon | Pop | Pav |
| E. Summer '04 | Whitefish | 1.22±0.01(34) AB | 1.07±0.02(31) C | 1.30±0.04(33) A | 1.09±0.04(9) BC | - | - |
| L. Summer '04 | | 1.12±0.01(24) AB | 0.98±0.02(26) C | 1.14±0.02(15) A | 1.07±0.01(15) B | - | - |
| Spring '05 | | | 1.11±0.02(11) B | 1.19±0.03(18) A | | 1.09±0.01(12) B | - |
| L. Summer '05 | | | 1.13±0.04(18) A | 0.99±0.02(15) B | 1.06±0.03(12) AB | | - |
| E. Summer '04 | | | - | 1.18±0.03(57) B | 1.16±0.03(37) B | 1.06±0.02(30) B | 1.27±0.02(21) A |
| L. Summer '04 | Sucker | - | 1.04±0.04(25) A | 1.13±0.03(11) A | 1.36±0.22(28) A | 1.11±0.01(66) A | 1.12±0.02(42) A |
| Spring '05 | | | 1.10±0.01(27) AB | 1.15±0.14(22) A | | 1.07±0.2(20) B | 0.99±0.08(14) AB |
| L. Summer '05 | | | - | 1.11±0.02(19) A | 1.06±0.02(20) A | | 1.08±0.02(18) A |

Letters represent significant differences among sites within a sampling period (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Condition factor = weight*100/length³.

- species not caught during sampling period

mykiss), cutthroat trout (*Oncorhynchus clarki*) and mountain whitefish are present in the upper reaches of the river. In the lower reaches cool-water fish such as suckers (mountain, white, and longnose), mooneye (*Hiodon tergisus* Lesueur), goldeye (*Hiodon alosoides*), and northern pike (*Esox lucius* Linnaeus) dominate. In between these extremes, we find both cold- and cool-water fish sharing the same habitat. The pressure of agriculture also increases in an eastward direction, and linked to these activities is an increase in water concentrations of herbicides, such as 2,4-D and MCPP (Mecoprop) and other compounds, such as cholesterol and desmosterol. OPs and CBs were not detected in the Oldman River in the present study; they have a short half-life and are usually only detected in water for a few days after application (Morrison & Wells 1981, Fulton & Key 2001). Furthermore, only a limited number of OPs and CBs were measured by Alberta Environment, due to limited resources. However, the effects, including AChE inhibition, of OPs and CBs can be detected even after a recovery period (Beauvais et al. 2000) and trace amounts of many different OPs and CB can have an affect on organisms because they act through a similar mechanism (Grue et al. 1991, Scholz et al. 2006).

OPs and CBs inhibit AChE, the enzyme that is responsible for the removal of the neurotransmitter acetylcholine, from synaptic clefts and motor endplates (Fulton & Key 2001). Measuring levels of AChE is an easy and inexpensive way of estimating exposure to OPs and CBs. Although there was no detectable gradient of OPs or CBs in the Oldman River, we did observe a species difference in AChE: whitefish always had lower levels of AChE as compared to suckers. The difference in AChE activity between the species could be due to a fundamental physiological difference. Davies et al. (1994)

detected different AChE activities in rainbow trout and four species of Australian fish exposed to pesticides Acephate and Fenitrothion. Van Dolah et al. (1997) reported that Mummichog (*Fundulus heteroclitus*) had lower levels of brain AChE compared to Red Drum (*Sciaenops ocellatus*) when exposed to the same concentrations of Azinphosmethyl. The lower activities of AChE in whitefish could also indicate that whitefish are exposed to higher concentrations of OPs and/or CBs, either through differences in microhabitat increasing exposure to waterborne pesticides, or a difference in pesticide loading through diet. Whitefish feed on dipteran, ephemeropteran, trichopteran and some terrestrial invertebrates (Pontius & Parker 1973, Thompson & Davies 1976, Ellison 1980), while the main food source for suckers is detritus (Saint-Jacques et al. 2000), ostracods, chironomids and amphipods (Johnson & Dropkin 1995).

Along with differences in AChE activities related to fish species, small but significant site-related differences were also observed. The lower AChE activities in whitefish in spring 2005 at easterly sites may be due to an increase in pesticide exposure in early summer, a period of intensive pesticide application in Southern Alberta. Most OPs and CBs have the same mode of action, so while water concentrations of individual pesticides may be at trace amounts and not affect fish, small amounts of many different pesticides acting together may inhibit AChE (Grue et al. 1991). In late summer 2005 (late July to early August), AChE was however lower in western sites for both species. The lower AChE activity at western cooler sites could potentially be due to a temperature effect: Hogen (1970) described a linear relationship between AChE and temperature in bluegills. The relationship between temperature and pesticide exposure on AChE inhibition needs to be explored in greater depth, so that this marker of exposure can be

validated for use in a wide range of fish species and temperature regimes. Furthermore, AChE profile in mountain whitefish and suckers needs to be further characterized. Ours is a first set of data for AChE in whitefish and first seasonal data for sucker. Therefore, at present, it can not be concluded with confidence that the AChE values detected in this study are an indication of inhibition of AChE or if these are normal values.

2.4.2 *Physiological Stress Response*

The present study also provided a first set of data on plasma cortisol in mountain whitefish. Cortisol is a steroid hormone that plays a major role in maintaining energy stores and energy mobilization at times of stress in fish (Hontela 1998, Mommsen et al. 1999, Schreck et al. 2001, Barton 2002). Temperature is a known stressor and can increase plasma cortisol in fish at water temperatures outside of the thermal preference (Davis 2004). Plasma cortisol levels at sites outside of the whitefish thermal preference range were higher in whitefish than suckers; whitefish sampled at Pearce Rd in early summer 2004 had very high plasma cortisol levels, caused most likely by the high water temperature (23-25°C) on days of sampling. Interestingly, suckers had higher plasma cortisol than whitefish at Ft. MacLeod, the most western and also coldest site where suckers are present in abundance in the Oldman River. Temperature below the optimal temperature range is known in some species to increase plasma cortisol in fish, as temperatures above this range do. Weber & Bosworth (2005) found that channel catfish (*Ictalurus punctatus*) reared at cold temperatures had higher cortisol levels than their controls. However, other studies in striped bass (*Morone saxatilis*) and sunshine bass (*Morone chrysops* X *Morone saxatilis*) have shown that cold water temperature had no effect or lowered plasma cortisol (Davis & Parker 1990, Davis & Peterson 2006).

Several studies have shown, however, that plasma cortisol levels are species-dependent; two species may have different levels of plasma cortisol when exposed to the same stressor (Lacroix & Hontela 2004, Jentoft et al. 2005).

It is difficult to differentiate between the acute and the chronic effects of stressors on plasma cortisol in field studies. Plasma cortisol levels increased in an eastward direction for whitefish in late summer 2004 and during both sampling periods in 2005, while suckers followed the same pattern in 2005. This increase in plasma cortisol follows both the pesticides/chemical gradient and temperature gradient, suggesting that one or both of these factors act as stressors on both whitefish and suckers. However, there is also evidence in the literature to suggest that pesticides impair cortisol secretion in fish chronically exposed to low concentrations (Dorval et al. 2005, Thangavel et al. 2005). Dimethoate, an organophosphate, is known to inhibit steroid hormone production by blocking the steroidogenic acute regulatory (StAR) protein (Walsh et al. 2000). Plasma cortisol levels at all sites for suckers were within the range found for suckers at agriculturally impacted sites and well below the normal values for suckers (Dorval et al. 2005). Thus, our data suggest that the stress response in suckers in the Oldman River is impaired, possibly by agricultural chemicals. Normal plasma cortisol values are not known for mountain whitefish, since this species has not been investigated previously. The cortisol and AChE data suggest that whitefish may be highly vulnerable to the combined stress effects of pesticides and increased temperature. Further studies are needed to determine what the combined effects of pesticides and temperature are in order to determine if cortisol levels in this study were impaired by pesticides.

Liver glycogen is a stored energy source that gets catabolised into a useable energy source, glucose, during times of stress (Wendelaar Bonga 1997). Whitefish had lower levels of liver glycogen compared to suckers at all sites and sampling periods. Our study is a first comparative study to provide data on liver glycogen in these two fish species, thus, as with AChE activities and plasma cortisol, it is difficult to characterize the effects of stressors and the fundamental differences between species. The lower liver glycogen in whitefish could be due to 1) a difference in sensitivity to temperature and/or pesticides, indicating that whitefish show greater signs of stress, or 2) a physiological difference between the two species and a lower capacity for storage of glycogen in the liver of whitefish. Studies that compared liver glycogen in different species have shown that there are species-specific differences in liver glycogen levels (Krogdahl et al. 2004, Tan et al. 2006).

Site differences in liver glycogen reserves were also detected in the present study. Whitefish had lower liver glycogen levels, and very high cortisol levels, at Pearce Rd in early summer 2004, most likely associated with the high temperature during sampling. Liver glycogen reserves in both whitefish and suckers decreased along a west-east gradient, suggesting that the increase of temperature and/or pesticides is recognized as a stressor, leading to a depletion of glycogen reserves. Liver glycogen levels have been shown to decrease when fish are exposed to increased temperature or pesticides; matrinxã (*Brycon cephalus*) exposed to the organophosphate, Folidol 600 (methyl parathion) (de Aguiar et al. 2004) and steelhead trout exposed to increased water temperature both showed a decrease in liver glycogen (Viant et al. 2003).

One of the characteristic responses associated with an increase in plasma cortisol is a rise in plasma glucose, the energy source used to maintain homeostasis (Hontela 1998, Mommsen et al. 1999). Warm water temperature has been found to increase plasma glucose in eelpout (Van Dijk et al. 1999) and other fish. Plasma glucose levels in both whitefish and suckers were similar and were not influenced by site, even at Pearce Rd in summer 2004 where plasma cortisol was very high in whitefish. These results suggest that both fish species were able to maintain their glycaemia, despite other physiological signs of stress (increased plasma cortisol and decreased liver glycogen). Thus, plasma glucose was not a sensitive physiological indicator of stress in fish in our field study.

Condition factor, often used as an indicator of overall health, was not affected in whitefish or suckers sampled in this study. Again, it is at present difficult to conclude whether the condition factors were within the normal range, especially for whitefish, because this species has not been investigated in great detail. One study reported that mountain whitefish at a reference site had a condition factor of 1.14 ± 0.22 while whitefish exposed to bleached kraft mill effluent had a condition factor of 1.20 ± 0.18 and 1.07 ± 0.17 (Swanson et al. 1994). Studies measuring the condition factor in suckers are numerous but condition factor for suckers in reference systems can range from 1.10 to 1.82 while condition factor for suckers in contaminated sites range from 1.23 to 1.72 (Munkittrick & Dixon 1989, Swanson et al. 1994). It is known however that if water temperatures continue to rise, foraging frequency and durations will decrease (Neill & Magnuson 1974), which would lead to a reduction in available energy, and over time would lead to a decreased condition factor.

Global climate change and increased use of pesticides are linked because with increasing air temperatures, new pests and diseases affecting both agriculture and human health are able to survive in areas that were once too cold (Bloomfield et al. 2006). The ability to control these pests usually relies on the use of pesticides (Bloomfield et al. 2006). A study of the impact that global climate change will have on agriculture in Canada and the United Kingdom shows that weeds are likely to evolve rapidly with climate change and that milder winters will increase the survival of some pests and diseases, and allow their establishment in new areas (Smith & Almaraz 2004, Bloomfield et al. 2006). Changes in temperature will also lead to a shift in crop type and with these new crops come their associated pests (Smith & Almaraz 2004, Bloomfield et al. 2006). Some areas will experience an increase in arable land while other areas will need irrigation to adequately supply water to fields, increasing water withdrawals (Bloomfield et al. 2006). Increased water withdrawals will most likely result in further water temperature increases and greater evaporation in rivers used for irrigation (Rosenberg et al. 1999). Furthermore with less water and continued input of pesticides into aquatic systems fish may be exposed to higher concentrations of pesticides.

Global climate change models suggest as water temperatures continue to increase there may also be a shift in the distribution of fish with decreases in cold-water species and the invasion of cool- or warm-water species (De Stasio et al. 1996, Eaton & Scheller 1996). At present, cold-water fish in the Oldman River are protected from the movement of some cool-water fish into the upper reaches of the river due to cold water temperatures and weirs. However, during high waters and/or as a result of improvements to fish ladders, cold-water fish (such as whitefish) might be extirpated from the Oldman River if

water temperatures continue to increase. Daufresne et al. (2003) have been monitoring water temperature increases in the Upper Rhône River in France and they observed a shift in community structure with displacement of cold-water fish and an increase in warm-water fish. Furthermore, studies on zooplankton show that high temperatures and exposure to toxicants lead to smaller bodied organisms which could affect the feeding in fish and lead to changes in fish distribution based on food availability (Moore & Folt 1993).

2.5 Conclusion

In conclusion, species-specific differences in AChE activities and responses of the physiological stress axis were detected in whitefish and suckers, suggesting that whitefish are a more sensitive species to temperature and pesticide stress. Whitefish appear to be at greater risk if water temperatures and pesticides inputs continue to increase. Further monitoring of the Oldman River and a greater understanding of mountain whitefish physiology are needed to determine if whitefish populations are healthy and stable. Studies are also needed to understand the interactions of increased temperature and pesticide exposure on the organisms within the impacted aquatic systems.

2.6 References

- Altinok, I., Capkin, E., Karahan, S., & Boran, M. 2006. Effects of water quality and fish size on toxicity of methiocarb, a carbamate pesticide, to rainbow trout. *Environmental Toxicology and Pharmacology* 22:20-26.
- Beauvais, S.L. Jones, S.B., Brewer, S.K., & Little, E.E. 2000. Physiological measures of neurotoxicity of diazinon and malathion to larval rainbow trout (*Oncorhynchus mykiss*) and their correlation with behavioural measures. *Environmental Toxicology and Chemistry* 19:1875-1880.
- Bartell, S.M., Breck, J.E., Gardner, R.H., & Brenkert, A.L. 1986. Individual parameter perturbation and error analysis of fish bioenergetics models. *Canadian Journal of Fisheries and Aquatic Science* 43:160-168.
- Barton, B.A. 2002. Stress in fishes: A diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and Comparative Biology* 42:517-525.
- Beitinger, T.L. & Magnuson, J.J. 1975. Influence of social rank and size on thermoselection behaviour of bluegill (*Lepomis macrochirus*). *Journal of the Fisheries Research Board of Canada* 32:2133-2136.
- Bhattacharya, S. 1993. Target and non-target effects of anticholinesterase pesticides in fish in: Proceeding of the second European conference on toxicology. *Science of the Total Environment Supplement*:859-866.
- Bleau, H., Daniel, C., Chevalier, G., van Tra, H., & Hontela, A. 1996. Effects of acute exposure to mercury chloride and methylmercury on plasma cortisol, T3, T4, glucose, and liver glycogen in rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology* 34:221-245.
- Bloomfield, J.P., Williams, R.J., Goody, D.C., Cape, J.N., & Guha, P. 2006. Impacts of climate change on the fate and behaviour of pesticides in surface and groundwater – a UK perspective. *Science of the total Environment* 369:163-177.
- Brett, J.R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). *American Zoologist* 11:99-113.
- Brewer, S.K., Little, E.E., DeLonay, A.J. Beauvais, S.L., Jones, S.B., & Ellersieck, M.R. 2001. Behavioral dysfunctions correlate to altered physiology in rainbow trout (*Oncorhynchus mykiss*) exposed to cholinesterase-inhibiting chemicals. *Archives of Environmental Contamination and Toxicology* 40:70-76.

- Capkin, E., Altinok, I., & Karahan, S. 2006. Water quality and fish size affect toxicity of endosulfan, an organochlorine pesticide, to rainbow trout. *Chemosphere* 64:1793-1800.
- Chuiko, G.M. 2000. Comparative study of acetylcholinesterase and butyrylcholinesterase in brain and serum of several freshwater fish: specific activities and in vitro inhibition by DDVP, an organophosphorus pesticide. *Comparative Biochemistry and Physiology* 127C:233-242.
- Cormier, S.M., Smith, M., Norton, S., & Neiheisel, T. 2000. Assessing ecological risk in watersheds: A case study of problem formulation in the Big Darby Creek Watershed, Ohio, USA. *Environmental Toxicology and Chemistry* 19:1082-1096.
- Daufresne, M., Roger, M.C., Capra, H., & Lamouroux, N. 2003. Long-term changes within the invertebrate and fish communities of the Upper Rhône River: effects of climatic factors. *Global Change Biology* 10:124-140.
- Davies, P.E., Cook, L.S.J., & Goenarso, D. 1994. Sublethal responses to pesticides of several species of Australian freshwater fish and crustaceans and rainbow trout. *Environmental Toxicology and Chemistry* 13:1341-1354.
- Davis, K.B. 2004. Temperature affects physiological stress responses to acute confinement in sunshine bass (*Morone chrysops* X *Morone saxatilis*). *Comparative Biochemistry and Physiology* 139A:433-440.
- Davis, K.B. & Parker, N.C. 1990. Physiological stress in striped bass: effect of acclimation temperature. *Aquaculture* 91:349-358.
- Davis, K.B. & Peterson, B.C. 2006. The effect of temperature, stress, and cortisol on plasma IFG-I and IGFBPs in sunshine bass. *General and Comparative Endocrinology* 149:219-225.
- de Aguiar, L.H. Moraes, G. Avilez, I.M., Altran, A.E., & Corrêa, C.F. 2004. Metabolical effects of Folidol 600 on the neotropical fish matrinxã, *Brycon cephalus*. *Environmental Research* 95:224-230.
- De Stasio, B.T., Jr., Hill, D.K., Kleinhans, J.M., Nibbelink, N.P., & Magnuson, J.J. 1996. Potential effects of global climate change on small north-temperate lakes: Physics, fish, and plankton. *Limnology and Oceanography* 41:1136-1149.
- Dill, L.M. 1987. Animal decision making and its ecological consequences: the future of aquatic ecology and behaviour. *Canadian Journal of Zoology* 65:803-811.
- Dorval, J., Leblond, V., Deblois, C., & Hontela, A. 2005. Oxidative stress and endocrine endpoints in white sucker (*Catostomus commersoni*) from a river impacted by agricultural chemicals. *Environmental Toxicology and Chemistry* 24:1273-1280.

- Eaton, J.G., McCormick, J.H., Goodno, B.E., O'Brien, D.G., Stefany, H.G., Hondzo, M., & Scheller, R.M. 1995. A field information-based system for estimating fish temperature tolerances. *Fisheries* 20:10-18.
- Eaton, J.G. & Scheller, R.M. 1996. Effects of climate warming on fish thermal habitats in streams of the United States. *Limnology and Oceanography* 41:1109-1115.
- Ellison, J.P. 1980. Diets of mountain whitefish, *Prosopium williamsoni* (Girard), and brook trout, *Salvelinus fontinalis* (Mitchill), in the Little Walker River, Mono County, California. *California Fish and Game* 66:96-104.
- Ellman, G.L., Courtney, K.D., Anders, V., Jr., & Featherstone, R.M. 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 7:88-95.
- Frank, R. & Sirons, G.J. 1979. Atrazine: Its use in corn production and its loss to stream waters in southern Ontario, 1975-1977. *The Science of the Total Environment* 12:223-239.
- Fulton, M.H. & Key, P.B. 2001. Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticide exposure and effects. *Environmental Toxicology and Chemistry* 20:37-45.
- Gordon, A.S., Gordon, E.P., & Dyer, B.J. 2006. Susceptibility of two fishes (*Oreochromis niloticus* and *Cyprinodon variegatus*) to *Pfiesteria shumwayae* and its associated toxin: Influence of salinity. *Harmful Algae* 5:542-547.
- Grue, C.E., Hart, A.D.M., & Mineau, P. 1991. Biological consequences of depressed brain cholinesterase activity in wildlife, In: *Cholinesterase-inhibiting insecticides* ed. Mineau. P. Elsevier. Amsterdam.
- Hasspieler, B.M., Behar, J.V., Carlson, D.V., Watson, D.E., & Di Giulio, R.T. 1994. Susceptibility of channel catfish (*Ictalurus punctatus*) and brown bullhead (*Ameiurus nebulosa*) to oxidative stress: a comparative study. *Aquatic Toxicology* 28:53-64.
- Hogan, J.W. 1970. Water temperature as a source of variation in specific activity of brain acetylcholinesterase of bluegills. *Bulletin of Environmental Contamination & Toxicology* 5:347-353.
- Hontela, A. 1997. Endocrine and physiological responses of fish to xenobiotics: Role of glucocorticosteroid hormones. *Reviews in Toxicology* 1:1-46.
- Hontela, A. 1998. Interrenal dysfunction in fish from contaminated sites: In vivo and in vitro assessment. *Environmental Toxicology and Chemistry* 17:44-48.

- Hontela, A. Dumont, P., Duclos, D., & Fortin, R. 1995. Endocrine and metabolic dysfunction in yellow perch, *Perca flavescens*, exposed to organic contaminants and heavy metals in the St. Lawrence River. *Environmental Toxicology and Chemistry* 14:725-731.
- Ihnat, J.M. & Bulkley, R.V. 1984. Influence of acclimation temperature and season on acute temperature preference of adult mountain whitefish, *Prosopium williamsoni*. *Environmental Biology of Fishes* 11:29-40.
- Jeffries, K.M. 2007. Endocrine disrupting compounds and responses of longnose dace in the South Saskatchewan River Basin. Masters Thesis, University of Calgary, Calgary, Alberta.
- Jentoft, S., Aastveit, A.H., Torjesen, P.A., & Andersen, Ø. 2005. Effects of stress on growth, cortisol, and glucose levels in non-domesticated Eurasian perch (*Perca fluviatilis*) and domesticated rainbow trout (*Oncorhynchus mykiss*). *Comparative Biochemistry and Physiology* 141A:353-358.
- Johnson, J.H. & Dropkin, D.S. 1995. Diel feeding chronology of six fish species in the Juniata River, Pennsylvania. *Journal of Freshwater Ecology* 10:11-18.
- Kuo, C.M. & Hsieh, S.L. 2006. Comparisons of physiological and biochemical responses between milkfish (*Chanos chanos*) and grass carp (*Ctenopharyngodon idella*) to cold shock. *Aquaculture* 251:526-536.
- Krogdahl, Å., Sundby, A., & Olli, J.J. 2004. Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*) digest and metabolize nutrients differently. Effects of water salinity and dietary starch level. *Aquaculture* 229:335-360.
- Lacroix, A. & Hontela, A. 2004. A comparative assessment of adrenotoxic effects of cadmium in two teleost species, rainbow trout, *Oncorhynchus mykiss*, and yellow perch, *Perca flavescens*. *Aquatic Toxicology* 67:13-21.
- Laflamme, J.-S., Couillard, Y., Campbell, P.G.C., & Hontela, A. 2000. Interrenal metallothionein and cortisol secretion in relation to Cd, Cu, and Zn exposure in yellow perch, *Perca flavescens*, from Abitibi lakes. *Canadian Journal of Fisheries and Aquatic Science* 57:1692-1700.
- Lappivaara, J. & Marttien, S. 2005. Effects of iron overload and simulated winter conditions on acute physiological stress response of whitefish, *Coregonus lavaretus*. *Ecotoxicology and Environmental Safety* 60:157-168.
- Linde, A.R., Sánchez-Galán, S., Klein, D., García-Vázquez, E., & Summer, K.H. 1999. Metallothionein and heavy metals in brown trout (*Salmo trutta*) and European eel

- (*Anguilla anguilla*): A comparative study. *Ecotoxicology and Environmental Safety* 44: 168-173.
- Moore, M. & Folt, C. 1993. Zooplankton body size and community structure: Effects of thermal and toxicant stress. *Trends in Ecology and Evolution* 8:178-183.
- Morrison, B.R.S. & Wells, D.E. 1981. The fate of fenitrothion in a stream environment and its effect on the fauna, following aerial spraying of a Scottish forest. *The Science of the Total Environment* 19:233-252.
- Mommsen, T.P., Vijayan, M.M., & Moon, T.W. 1999. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. *Reviews in Fish Biology and Fisheries* 9:211-268.
- Munkittrick, K.R. & Dixon, D.G. 1988. Growth, fecundity, and energy stores of white sucker (*Catostomus commersoni*) from lakes containing elevated levels of copper and zinc. *Canadian Journal of Fisheries and Aquatic Science* 45:1355-1365.
- Neill, W.H. & Magnuson, J.J. 1974. Distributional ecology and behavioral thermoregulation of fishes in relation to heated effluent from a power plant at Lake Monona, Wisconsin. *Transactions of the American Fisheries Society* 103:663-709.
- Oldman Watershed Counsel. 2006. Oldman river basin map. www.oldmanbasin.org/images/basin_sm.jpg. Accessed February 2007.
- Pavlov, D.D., Chuiko, G.M. Gerassimov, Y.V. & Tonkopyi, V.D. 1992. Feeding behavior and brain acetylcholinesterase activity in bream (*Abramis brama* L.) as affected by DDVP, an organophosphorus insecticide. *Comparative Biochemistry and Physiology* 103C:563-568.
- Pontius, R.W. & Parker, M. 1973. Food habits on the mountain whitefish *Prosopium williamsoni* (Girard). *Transactions of the American Fisheries Society* 102:764-773.
- Rosenberg, N.J., Epstein, D.J., Wang, D., Vail, L. Srinivasan, R., & Arnold, J.G. 1999. Possible impacts of global warming on the hydrology of the Ogallala aquifer region. *Climatic Change* 42:677-692.
- Saint-Jacques, N., Harvey, H.H., & Jackson, D.A. 2000. Selective foraging in the white sucker (*Catostomus commersoni*). *Canadian Journal of Zoology* 78:1320-1331.
- Samuel, M. & Sastry, K.V. 1989. *In vivo* effect of monocrotophos on the carbohydrate metabolism of the freshwater snake head fish, *Channa punctatus*. *Pesticide Biochemistry and Physiology* 34:1-8.

- Sandahl, J.F., Baldwin, D.H., Jenkins, J.J., & Scholz, N.L. 2005. Comparative thresholds for acetylcholinesterase inhibition and behavioral impairment in coho salmon exposed to chlorpyrifos. *Environmental Toxicology and Chemistry* 24:136-145.
- Schindler, D.W., Bayley, S.E., Parker, B.R., Beaty, K.G., Cruikshank, D.R., Fee, E.J., Schindler, E.U., & Stainton, M.P. 1996. The effects of climatic warming on the properties of boreal lakes and streams at the experimental lakes area, Northwestern Ontario. 41:1004-1017.
- Schreck, C.B., Contreras-Sanchez, W., & Fitzpatrick, M.S. 2001. Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture* 197:3-24.
- Scholz, N.L., Truelove, N.K., Labenia, J.S., Baldwin, D.H., & Collier, T.K. 2006. Dose-additive inhibition of Chinook salmon acetylcholinesterase activity by mixtures of organophosphate and carbamate insecticides. *Environmental Toxicology and Chemistry* 25:1200-1207.
- Smith, D.L. & Almaraz, J.J. 2004. Climate change and crop production: contributions, impacts, and adaptations. *Canadian Journal of Plant Pathology* 26:253-266.
- Stefan, H.G., Hondzo, M., & Fang, X. 1993. Lake water quality modeling for projected future climate scenarios. *Journal of Environmental Quality* 22:417-431.
- Sullivan, K., Martin, D.J., Cardwell, R.D., Toll, J.E., & Duke, S. 2000. An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria. Sustainable Ecosystems Institute. Portland, Oregon.
- Swanson, S.M., Schryer, R., Shelast, R., Kloepper-Sams, P.J., & Owens, J.W. 1994. Exposure of fish to biologically treated bleached-kraft mill effluent. 3. Fish habitat and population assessment. *Environmental Toxicology and Chemistry* 13:1497-1507.
- Tan, Q., Xie, S., Zhu, X., Lei, W., & Yang, Y. 2006. Effect of dietary carbohydrate sources on growth performance and utilization for gibel carp (*Carassius auratus gibelio*) and Chinese longsnout catfish (*Leiocassis longirostris* Günther). *Aquaculture Nutrition* 12:61-70.
- Teles, M., Pacheco, M., & Santos, M.A. 2007. Endocrine and metabolic responses of *Anguilla Anguilla* L. caged in a freshwater-wetland (Pateira de Fermentelos-Portugal). *Science of the Total Environment* 372:562-570.
- Thangavel, P., Sumathiral, K., Karthikeyan, S., & Ramaswamy, M. 2005. Endocrine response of the freshwater teleost, *Sarotherodon mossambicus* (Peters) to dimecron exposure. *Chemosphere* 61:1083-1092.

- Thompson, G.E. & Davies, R.W. 1976. Observations on the age, growth, reproduction, and feeding of mountain whitefish (*Prosopium williamsoni*) in the Sheep River, Alberta. Transactions of the American Fisheries Society 105:208-219.
- Van Dijk, P.L.M., Tesch, C., Hardewig, I., & Pörtner, H.O. 1999. Physiological disturbances at critically high temperatures: a comparison between stenothermal antarctic and eurythermal temperate eelpouts (*Zoarctidae*). The Journal of Experimental Biology 202:3611-3621.
- Van Dolah, R.F., Maier, P.P., Fulton, M.H., & Scott, G.I. 1997. Comparison of azinphosmethyl toxicology to juvenile red drum (*Sciaenops ocellatus*) and the mummichog (*Fundulus heteroclitus*). Environmental Toxicology and Chemistry 16:1488-1493.
- Viant, M.R., Werner, I., Rosenblum, E.S., Gantner, A.S., Tjeerdema, R.S., & Johnson, M.L. 2003. Correlation between heat-shock protein induction and reduced metabolic condition in juvenile steelhead trout (*Oncorhynchus mykiss*) chronically exposed to elevated temperature. Fish Physiology and Biochemistry 29:159-171.
- Walsh, L.P., Webster, D.R., & Stocco, D.M. 2000. Dimethoate inhibits steroidogenesis by disrupting transcription of the steroidogenic acute regulatory (StAR) gene. Journal of Endocrinology 167:253-263.
- Weber, T.E. & Bosworth, B.G. 2005. Effects of 28 day exposure to cold temperature or feed restriction on growth, body composition, and expression of genes related to muscle growth and metabolism in channel catfish. Aquaculture 246:483-492.
- Wendelaar Bonga, S.E. 1997. The stress response in fish. Physiological Reviews 77:591-625.

Chapter 3.

Effects of Dimethoate and temperature on acetylcholinesterase activity and cortisol secretion in rainbow trout (*Oncorhynchus mykiss*): *in Vivo* and *in Vitro* exposure

3.1 Introduction

Dimethoate (O,O-dimethyl-S-(N-methylcarbamoylmethyl) phosphorodithioate), an organophosphate (OP) used widely in agriculture for crop protection as a systemic and on-contact anticholinergic pesticide, targets a broad range of insects and mites (WHO 1989). It is used on a variety of fruit, vegetable and field crops, and also to control houseflies indoors (Farag et al. 2006). Dimethoate is a low persistence pesticide which has a degradation half-life of 4-16 days but can last longer depending on the conditions (Wauchope et al. 1992, Athanasopoulos et al. 2000). Alkalinity and temperature influence its degradation, with low temperature and low pH increasing the half-life (Athanasopoulos et al. 2000). Organophosphates such as Dimethoate are highly soluble in water and can leach into nearby water sources and affect aquatic organisms (Tuduri et al. 2006).

Exposure of non-target organisms such as vertebrates to Dimethoate affects behaviour (Dell’Omo et al. 1997, De Mel & Pathiratne 2005, Farag et al. 2006), reproductive function (Begum & Vijayaraghavan 1995, Farag et al. 2006) and muscle glycogen levels (Begum & Vijayaraghavan 1999). Dimethoate also causes a reduction in food consumption and body weight in mammals (Farag et al. 2006), and damage to gill, kidney, and liver tissues in fish (Gill et al. 1988). Exposure to Dimethoate inhibits acetylcholinesterase (AChE) in the brain (Dell’Omo et al. 1997, Frasco & Guilhermino 2002, De Mel & Pathiratne 2005, Farag et al. 2006).

Exposure to stressors, including chemical stressors, such as pesticides, and physical stressors, such as suboptimal water temperature, can activate the physiological stress response, a suite of responses that facilitate maintenance of homeostasis (Wendelaar Bonga 1997, Schreck et al. 2001). Plasma cortisol and glucose levels, or liver glycogen, are physiological indicators used to determine if fish are stressed and if the hypothalamo-pituitary-interrenal axis is functioning normally (Hontela 1998). Acute exposures to pesticides elevate plasma cortisol (Teles et al. 2007) while chronic exposure may actually impair cortisol synthesis in fish species (Dorval et al. 2005, Thangavel et al. 2005). Although the effects of Dimethoate on cortisol synthesis have not been investigated in fish or other vertebrates, Dimethoate has been shown to block the steroidogenic pathway, specifically the StAR protein required for cholesterol uptake by the mitochondrion, in the testis of rats and inhibit synthesis of testosterone (Walsh et al. 2000). Since synthesis of cortisol, another steroid hormone, also relies on StAR, exposure to Dimethoate could block cortisol synthesis in the steroidogenic interrenal tissue in teleosts.

Cortisol synthesis and plasma cortisol levels in fish can be also influenced by temperature. When fish are exposed to temperatures outside their thermal preference range (both above and below); there is an increase in plasma cortisol levels (Davis & Parker 1990, Davis 2004, Davis & Peterson 2006). Suboptimal temperatures can also have negative effects on growth rate and reproduction (Davis & Parker 1990, Martikainen & Rantalainen 1999, Davis 2004, Davis & Peterson 2006). Temperature can not only affect the physiology and behaviour of organisms, but also the fate of chemicals in the environment and their toxicity (Martikainen & Rantalainen 1999). Chemicals that

cause oxygen deficiency or increase metabolic activity tend to be more toxic at higher temperatures (Martikainen & Rantalainen 1999) whereas low temperatures can increase toxicity by increasing exposure duration (Athanasopoulos et al. 2000). The effects of temperature on OP pesticides exposure and the inhibition of AChE vary considerably; some studies (Beauvais et al. 2002, Phillips et al. 2002) reported that temperature does not affect AChE inhibition while others (Hogan 1970, Zinkl et al. 1987) showed a linear relationship between water temperature and AChE inhibition. With the threat of global climate change, it is important to understand the effect that increasing temperatures will have on aquatic organisms and their exposure to chemicals such as pesticides. Pesticide use is expected to increase with increasing global temperature, to control new pests and expanding pest populations. Thus within the climate change scenario, aquatic organisms may be potentially exposed to both increased temperatures and increased pesticide inputs (Bloomfield et al. 2006).

*The objectives of this study were to investigate the effects of Dimethoate and temperature on cortisol secretion and the physiological stress response, and AChE activity in rainbow trout (*Oncorhynchus mykiss*). It is hypothesised that exposure to Dimethoate will inhibit cortisol secretion and AChE activity in all tissues, and higher temperature will increase AChE activity and the toxicity of Dimethoate.*

3.2 Materials and Methods

a. Test Species

Immature male and female rainbow trout, *O. mykiss*, obtained from Allison Creek Brood Trout Station (Coleman, Alberta, Canada), were transferred on arrival into a

maintenance tank (600L) for a 2-week acclimation period (12L:12D, 15±1°C, oxygen saturated water, pH 7.8). Fish were fed commercial trout food (Silver Cup salmon crumbles, Nelson & Sons) once a day.

3.2.1 In Vivo exposure to waterborne Dimethoate (pure grade)

Following the acclimation period, fish (size matched groups of 10) were transferred into 6 experimental tanks (semi-static system, with a water renewal of 20% daily). All the tanks were located in the aquatic room maintained at air temperature of 11°C; each experimental tank had a submerged heater (Thermal Compact Pre-Set Heater).

a. Experimental treatments

Fish (average body weight 24.7±0.5 g, average body length 12.4±0.1 cm) were acclimatized to the experimental tanks (50 L) for one week, then they were exposed to 0% (control), 0.5%, or 1% LC₅₀ (LC₅₀ = 7.35 mg/L, Johnson and Finley 1980, WHO 1989) of pure grade Dimethoate (Riedel-de Haën 45449) dissolved in water, and either a cold or a warm water treatment for 21 days. The cold water tanks were maintained at 10±1°C for 21 days, using the air cooling system of the aquatic room. The warm water tanks were maintained at 19±1°C for 10 days using the submerged heaters. On Day 11 all the heaters were removed and for the next 10 days during the daily 20% water renewal, warm water (30±1°C) was added to the warm water treatment tanks at 0830h, to bring the water temperature up to 15±1°C. Water at 11°C was added to the cold water treatment tanks, to insure same level of disturbance in all the tanks. The Dimethoate concentrations were maintained at the experimental levels during water renewal. Water chemistry and temperature during the experiment are presented in Table 3.1 and Fig. 3.1.

Table 3.1. Temperature and water chemistry (mean \pm SEM) during Dimethoate (pure grade) and temperature treatments.

| Treatment | Temperature Day 1-10 (n=9) | Maximum Temperature Day 11-21 (n=12) | Dimethoate ^a mg/L (n=3) | pH (n=6) | Dissolved Oxygen (n=4) |
|----------------------------|----------------------------|--------------------------------------|------------------------------------|-----------------|------------------------|
| cold 0% LC ₅₀ | 10.44 \pm 0.13 | 10.17 \pm 0.07 | <0.1 | 7.63 \pm 0.07 | 9.98 \pm 0.77 |
| warm 0% LC ₅₀ | 20.11 \pm 0.18 | 15.58 \pm 0.17 | <0.1 | 7.64 \pm 0.09 | 8.28 \pm 0.77 |
| cold 0.5% LC ₅₀ | 9.72 \pm 0.12 | 9.75 \pm 0.13 | 31.6 \pm 0.3 | 7.97 \pm 0.06 | 9.83 \pm 0.81 |
| warm 0.5% LC ₅₀ | 18.61 \pm 0.27 | 15.08 \pm 0.17 | 29.6 \pm 2.3 | 7.92 \pm 0.09 | 6.87 \pm 0.36 |
| cold 1% LC ₅₀ | 10.89 \pm 0.11 | 11.13 \pm 0.07 | 110.2 \pm 28.2 | 7.80 \pm 0.05 | 7.01 \pm 0.41 |
| warm 1% LC ₅₀ | 17.44 \pm 0.31 | 15.13 \pm 0.11 | 78.0 \pm 18.9 | 7.86 \pm 0.06 | 6.25 \pm 0.19 |

^aWater samples were analyzed for Dimethoate by Alberta Research Council Laboratory (D. Humphries)




Figure 3.1. Hourly temperature profile for days 11-21 in the experimental treatment tanks.

b. Sampling

Fish were sampled between 0830h and 0930h to minimize the effects of diel hormonal variations. Fish were anaesthetized in a solution of 0.15 g/L of tricaine methanesulphonate (MS-222) and a blood sample was taken from the caudal vasculature with a 1 mL heparized syringe. Plasma was separated at 13000 rpm for 5 minutes and frozen in liquid nitrogen. The fish were euthanized by spinal transection, weights and lengths recorded and, the head kidneys were removed immediately and placed in minimal essential medium (MEM), liver, gills, muscle and brains were dissected and frozen in liquid nitrogen. Plasma and tissue samples were kept frozen at -80°C until analysis.

c. Acetylcholinesterase (AChE)

The AChE activity was measured with an assay modified from Ellman et al. (1961) and Chuiko (2000). Muscle samples were homogenized using Power Max Advanced Homogenizing System (VWR) (1 g of tissue to 3 mL phosphate buffer) and centrifuged at 5°C at 10000 rpm for 10 minutes. Brain samples were homogenized with a Pellet Pestle Motor (Kontes) (1g of tissue to 3 mL phosphate buffer) and centrifuged at 5°C at 13000 rpm for 3 minutes. A 10x dilution was used for both muscle and brain samples. Plasma, brain or muscle homogenate (2 µL) was pipetted into a microplate and 120 µL of ISO-OMPA (AChE only, Sigma Tetraisopropyl pyrophosphoramidate, T1505) or 120 µL of Tris Buffer (Total Cholinesterase: AChE and Butyrylcholinesterase (BChE), Tris Buffered Saline Tablets, Sigma T-5030) was added. The samples were incubated at room temperature for 10 minutes and 10 µL of DTNB (Sigma 5,5' Dithiobis (2-nitrobenzoic acid), D8130) and 10 µL Acetylthiochlorine Iodine (Sigma minimum 98% TLC, A5751) were added, followed by a 10 minute incubation period at room

temperature. The microplate was read at 405nm in a microplate reader every 2 minutes for 10 minutes. Concentration of AChE for each sample was measured from the slope; internal standards (Normal Serum Control, TC-TROL(N), Teco Diagnostics) and controls (1U/mL and 2 U/mL of eel acetylcholinesterase, Sigma C3389) were used to ensure the accuracy of the assay was maintained. Protein for muscle and brain samples was measured using the Bradford method (Bradford 1976).

d. Cortisol secretion (ACTH-challenge of head kidneys isolated from Dimethoate-exposed fish)

The head kidney was divided in two, half frozen for later analysis and the remaining portion used for the ACTH-challenge using the microplate method described by Benguira & Hontela (2000). The head kidney was cut into small fragments (1 mm³) and the coarse homogenate was divided into two portions (one for basal treatment and one for ACTH stimulation), pipetted into a 24 well microplate (1 mL of MEM/well). Following a 2 hrs incubation, the microplate was centrifuged and the supernatant removed. The basal treatment was then incubated in 1mL of MEM while the stimulated treatment was incubated in 1mL of MEM containing 2 IU of porcine adrenocorticotropin hormone (ACTH, Sigma Chemical Co A6303). After 1hr of incubation, the microplate was centrifuged and the supernatant was frozen for later analysis of cortisol by radioimmunoassay.

e. Physiological Stress Response

Concentrations of cortisol in the plasma were determined with diagnostic kits (MP Biomedicals Diagnostics Division 07-221102). Plasma glucose was determined with a spectrophotometer assay (512nm) using the GOD-PAP reagent (Roche 1929526).

Liver glycogen was measured with a method modified from Bleau et al. (1996). Liver samples were digested in KOH (1N) at room temperature; the sample was then precipitated with 1.5 N acetic acid and centrifuged at 13000 rpm for 10 minutes. Glycogen from the livers and standards (rabbit glycogen, Sigma, G8876-1G) was hydrolyzed to glucose with aminoglycosidase (Sigma A-7255) in a water bath at 37°C for 30 minutes. Glucose content was measured using the spectrophotometric assay described above.

f. Gill Na⁺/K⁺ ATPase activity

Na⁺/K⁺ -ATPase activity in the gills was determined using modified methods of Holliday (1985) and Morgan et al. (1998), as described in Levesque et al. (2003). In summary, 0.5g of gill tissue was homogenized in a phosphate buffer using a Pellet Pestle Motor (Kontes). Homogenates were added to 200 µL buffer A (NaCl 167 mM, KCl 50 mM, imidazole-HCl 33 mM, pH 7.2 with NaOH 1 M) or buffer B (NaCl 237 mM, ouabain 1.67 mM, imidazole-HCl 33 mM, pH 7.2 with NaOH 1 M). The Na⁺/K⁺ -ATPase activity was obtained by determining the difference in inorganic phosphate production between buffer A and buffer B using a spectrophotometer at 700nm. Protein was measured using the Bradford method (Bradford 1976). Enzyme activity was expressed as micromoles of inorganic phosphate per hour per milligram of protein.

3.2.2. In Vitro exposure of head kidney cells to Dimethoate

a. Preparation of head kidney cell suspensions

The method described by Leblond et al. (2001) was used for the head kidney cell preparations. Fish were anaesthetized with MS-222 (tricaine methanesulphonate), a blood sample was taken from the caudal vein and the fish were perfused through the heart with

a 0.7% saline and heparin solution to remove as much blood as possible. Head kidneys were then dissected out, placed in fresh MEM, cut into small pieces (<1 mm³) and placed into 2.5 ml MEM with 3.0 mg/mL DNAase (Sigma D8754) and 5.0 mg/mL collagenase/dispase (Boehringer Mannheim: 1-097-113). The cells were incubated at 22°C for 60 min with gentle agitation and cells were re-suspended every 15 min with a transfer pipette. After the digestion, the cells were agitated with a pipette to further break apart the tissue, then the cells were filtered with a 30 µm mesh cloth and centrifuged at 1000 rpm for 5 min at 15° C. The supernatant was removed, the pellet was re-suspended in MEM and cell density was adjusted to 75x10⁶ cells/mL. The cells were then distributed (75 µL of the cell suspension) in two 96 well microplates, preincubated for 120 min, one plate at 11±1° C and the other at 20±1° C.

b. Exposure to pure grade or commercial Dimethoate

Following the preincubation, the cells were centrifuged at 1000 rpm for 3 min, exposed to either pure grade Dimethoate (0 mg/L, 3 mg/L, 30 mg/L or 300 mg/L) (Riedel-de Haën 45449) or Cygon, the commercial dimethoate (0 mg/L, 3 mg/L, 30 mg/L, 60 mg/L, or 300 mg/L) in Ringer solution (269 mOsm/L), incubated for 60 min at either 11±1°C or 20±1°C, washed with 100 µL of Ringer solution, and stimulated with 1 IU/mL ACTH. Supernatant was collected for cortisol analysis.

Viability was determined using the trypan blue solution (0.4%, Sigma, T8154) and a counting chamber (VWR Scientific, 15370). Viability was expressed as a percentage the number of cells that took up the stain (dead) and those that did not (alive).

3.2.3 Statistical Analysis

For the *In Vivo* exposure experiments, a two-way factorial analysis (ANOVA) using JMP IN 5.1, with a post hoc Tukey-Kramer HSD test and a log transformation or Box Cox transformation for non-normal data, was used. Data that could not be transformed were analysed using a non-parametric Wilcoxon test with a Bonferoni correction. Linear regression with inverse prediction using JMP IN 5.1 was used to determine the LC₅₀ and EC₅₀ for *In Vitro* experiments. A paired t-test was used to determine if the LC₅₀ was different than the EC₅₀ and the difference between the two temperature treatments.

3.3 Results

3.3.1 *In Vivo* exposure to Dimethoate

a. AChE activity – a marker of pesticide exposure

The activity of AChE measured in plasma of control fish was lower than muscle and brain AChE (Table 3.2), with the highest activity detected in muscle ($p < 0.05$, ANOVA). The activity of AChE decreased in an exposure-dependent pattern in plasma, brain and muscle of rainbow trout exposed to Dimethoate through water, and the greatest inhibition was observed in muscle AChE (Table 3.2). BChE was not detected in any of the tissues (data not shown).

Table 3.2. AChE activity, mean±SE (n), in brain, muscle, and plasma in rainbow trout exposed to Dimethoate.

| Temperature treatment | Dimethoate (% LC ₅₀) | % AChE inhibition compared to controls | | |
|-----------------------|----------------------------------|--|-----------------|-----------------|
| | | Brain | Muscle | Plasma |
| cold | 0 | 0.00±0(10) E | 0.00±0 (10) E | 0.00±0 (10) E |
| | 0.5 | 18.89±5.2(10) D | 46.55±6.0(10) C | 27.65±4.9(10) D |
| | 1 | 57.22±3.2(9) BC | 87.91±2.4(9) A | 65.99±2.7(9) B |
| warm | 0 | 0.00±0 (8) e | 0.00±0 (8) e | 0.00±0 (8) e |
| | 0.5 | 2.46±1.5(7) e | 44.75±7.6(7) c | 25.03±7.4(7) d |
| | 1 | 42.22±3.5(9) cd | 86.12±3.8(9) a | 65.57±2.8(9) b |

Uppercase letters represent significant differences between brain, muscle, and plasma AChE inhibition for cold water, lowercase letters represent significant differences between brain, muscle, and plasma AChE inhibition in warm water (ANOVA and Tukey-Kramer HSD test $\alpha=0.05$).

Plasma AChE activity (Figure 3.2A) decreased with increasing exposure to Dimethoate, with significant differences ($p < 0.05$) between the control and the group exposed to 1% LC₅₀. The temperature treatments had no effect on plasma AChE activity at any concentration of Dimethoate.

Brain AChE activity was also inhibited with increasing exposure to Dimethoate (Figure 3.2B), again the effect was significant ($p < 0.05$) at the 1% LC₅₀ of Dimethoate. The warm water treatment decreased brain AChE activity in the control group (0% Dimethoate) but had no effect in the other exposure groups (0.5 and 1% LC₅₀).

Muscle AChE activity decreased significantly ($p < 0.05$) with increasing exposure to Dimethoate, in a concentration-dependant pattern, but there was no difference between the different temperature treatments (Figure 3.2C).

b. Physiological Stress Response

Plasma cortisol levels decreased with increasing concentrations of Dimethoate but the temperature treatments did not have an effect on cortisol (Figure 3.3A). Basal and ACTH-stimulated cortisol secretion of head kidneys isolated from fish exposed to Dimethoate in vivo was not different between the exposure groups or temperature treatment (Figure 3.3B). There were however significant differences in the variations between cortisol secretion at the different concentrations of Dimethoate (F-test for unequal variance).

There were no significant differences in plasma glucose or liver glycogen for either temperature treatment or Dimethoate concentrations ($p > 0.05$) (Table 3.3). Gill Na⁺/K⁺ ATPase activity increased with exposure to Dimethoate ($p < 0.05$) and warm temperature treatment increased the activity in the control group (Figure 3.4). There were




Figure 3.2. AChE activity in tissues of rainbow trout exposed to Dimethoate (pure grade) at two different temperatures. A) Plasma AChE, B) Brain AChE, C) Muscle AChE. Numbers in the bars indicate the number of fish in each treatment. Letters represent significant differences (Two-way ANOVA and Tukey-Kramer HSD test $\alpha=0.05$).




Figure 3.3. A) Plasma cortisol and (B) basal and ACTH-stimulated cortisol secretion (mean±SE) by isolated head kidneys of rainbow trout exposed to Dimethoate (pure grade) at two different temperatures. Numbers in the bars indicate the number of fish in each treatment. A) Upper-case letters represent significant differences (Wilxon non-parametric test with a bonferoni correction and Tukey-Kramer HSD test $\alpha=0.05$). B) Lower-case letters represent significant difference in basal cortisol while upper-case letters represent significant differences in stimulated cortisol (two-way ANOVA, with Tukey-Kramer HSD test $\alpha=0.05$).




Figure 3.4. Gill Na⁺/K⁺ ATPase activity (mean±SE) in rainbow trout exposed to Dimethoate (pure grade) at two different temperatures. Numbers in the bars indicate numbers of fish in each treatment. Capital letters represent significant differences (Two-way ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Data log transformed for statistical analysis.

Table 3.3. Plasma glucose, liver glycogen, condition factor and liver somatic index, mean ± SE(n), in rainbow trout exposed to Dimethoate (pure grade) at two different temperatures

| | Temperature | Dimethoate Treatment | | |
|------------------------|-------------|----------------------|-----------------------|---------------------|
| | | 0% LC ₅₀ | 0.5% LC ₅₀ | 1% LC ₅₀ |
| Plasma Glucose (mg/mL) | Cold Water | 0.87±0.04 (10) | 0.83±0.07 (10) | 0.85±0.11 (9) |
| | Warm Water | 0.87±0.07 (8) | 0.80±0.06 (7) | 0.84±0.06 (9) |
| Liver Glycogen (mg/g) | Cold Water | 63.73±9.3 (10) | 78.12±8.4 (10) | 61.59±9.5 (9) |
| | Warm Water | 88.83±11.6 (8) | 85.09±10.7 (7) | 83.14±14.8 (9) |
| Condition Factor | Cold Water | 1.30±0.19 (10) | 1.10±0.03 (10) | 1.11±0.01 (9) |
| | Warm Water | 1.12±0.04 (8) | 1.13±0.03 (7) | 1.13±0.02 (9) |
| Liver somatic index | Cold Water | 1.13±0.09 (10) | 1.33±0.07 (10) | 1.23±0.06 (9) |
| | Warm Water | 1.48±0.17 (8) | 1.42±0.15 (7) | 1.31±0.05 (9) |

Condition Factor: $\text{body weight} \times 100 / \text{length}^3$, Liver somatic index: $(\text{liver weight} / \text{body weight}) \times 100$ (Two-way ANOVA and Tukey-Kramer HSD test $\alpha=0.05$). Liver glycogen data log transformed for statistical analysis.

no significant differences for either temperature treatments or for Dimethoate concentrations ($p>0.05$) for condition factor or liver somatic index (LSI) (Table 3.3). Mortality occurred in all the warm water tanks, but was not dependent on Dimethoate concentration (two fish in control and 0.5% LC50 group, one fish in the 1% LC50 group). One fish in the cold water 0.5% LC50) dimethoate group died.



3.3.2 In Vitro exposure of head kidney cells to Dimethoate (pure grade)

a. Cortisol secretion

Pure grade Dimethoate did not have an effect on ACTH-stimulated cortisol secretion by isolated head kidney cells exposed to the pesticide in vitro at either $11\pm 1^\circ\text{C}$ ($R^2=0.04$, $P>0.05$) or $20\pm 1^\circ\text{C}$ ($R^2=0.33$, $P>0.05$) (Figure 3.5A). The EC_{50} (concentration that inhibits 50% of cortisol secretion) for pure Dimethoate could not be determined using inverse prediction because that level of inhibition was not observed. There was a significant difference ($p<0.05$) in cortisol secretion among the two temperatures, with higher cortisol levels detected at 20°C compared to 11°C (Figure 3.5B).

b. Viability

There was no change in viability with increasing concentrations of pure grade Dimethoate at 11°C ($R^2=0.04$, $P>0.05$, Figure 3.5C) but viability did decrease slightly but significantly ($R^2=0.33$, $P>0.05$) at 20°C (Figure 3.5C). Again, the LC_{50} (lethal concentration that kills 50% of cells) of pure Dimethoate could no be determined. Temperature did not have an effect on cell viability as there were no differences ($p>0.05$) in viability between 11°C and 20°C .






Figure 3.5. Cortisol secretion (ACTH-stimulated) and viability of head kidney cells exposed to Dimethoate (pure grade) in vitro at 11°C and 20°C. A) Cortisol secretion expressed as % Control at either 11°C or 20°C. Linear regression with inverse prediction was used to determine the slope of the line; solid black line is the regression for 11°C [$X=(Y+95.17)/0.02$], $R^2=0.04$, dashed line the regression for 20°C [$X=(Y+98.87)/0.04$], $R^2=0.33$, $n=11$ for 11°C and $n=13$ for 20°C, n indicating number of fish sampled, with individual wells pooled together. The solid grey line represents the predicted LC₅₀. B) Cortisol secretion expressed as % of Control at 20°C. C) Viability (expressed as % of Control) at either 11°C or 20°C, The solid line is the regression for 11°C [$X=(Y+86.54)/0.02$], $R^2=0.04$, and the dashed line is the regression for 20°C [$X=(Y+90.95)/0.08$], $R^2=0.33$. $n=8$ for all groups.



3.3.3 *In Vitro* exposure of head kidney cells to commercial dimethoate (Cygon)

a. *Cortisol secretion*

Cortisol secretion decreased ($p < 0.05$) with increasing concentrations of commercial dimethoate, Cygon, at both 11°C ($R^2 = 0.45$) and 20°C ($R^2 = 0.72$) (Figure 3.6A). The EC_{50} at 11°C, determined as 71.45 mg/L Cygon dimethoate (32.18-109.06 mg/L dimethoate 95% confidence limit), was significantly lower than the EC_{50} identified at 20°C as 118.10 mg/L Cygon dimethoate (97.24-141.67 mg/L Cygon dimethoate 95% confidence limit). Temperature had a significant effect ($p < 0.05$) on cortisol secretion, with higher cortisol secretion at 20°C than 11°C (Figure 3.6B).

b. *Viability*

There was a significant decrease ($p < 0.05$) in viability with increasing concentrations of Cygon dimethoate at both 11°C and 20°C (Figure 3.6C), however, temperature did not have a significant effect ($p > 0.05$) on cell viability. The LC_{50} for 11°C, determined as 101.14 mg/L Cygon dimethoate (73.45-131.70 mg/L Cygon dimethoate 95% confidence limit), was not significantly different from the LC_{50} at 20°C determined as 93.17 (66.97-120.90 mg/L Cygon dimethoate 95% confidence limit). At 11°C the LC_{50} was significantly greater than the EC_{50} however there was no difference between the EC_{50} and LC_{50} at 20°C.



C)




Figure 3.6. Cortisol secretion (ACTH-stimulated) and viability of head kidney cells exposed to Dimethoate (commercial grade) in vitro at 11°C and 20°C. A) Cortisol secretion expressed as % Control at either 11°C or 20°C. Linear regression with inverse prediction used to determine the slope of the line; solid black line is the regression for 11°C [$X=(Y+68.40)/0.26$], $R^2=0.45$, the dashed line is the regression for 20°C [$X=(Y+86.26)/0.31$], $R^2=0.72$. The grey line represents the EC_{50} for 11°C and 20°C: 71.45 mg/L dimethoate and 118.24 mg/L dimethoate respectively. $n=10$ for 11°C and 20°C, n indicating number of fish sampled, individual wells were pooled together. B) Cortisol secretion expressed as % of Control at 20°C, C) Viability (expressed as % of Control at either 11°C or 20°C, The black solid line is the regression for 11°C [$X=(Y+73.46)/0.23$] and the dashed black line is the regression for 20°C [$X=(Y+71.14)/0.23$]. The grey line represents the LC_{50} for 11°C and 20°C: 101.14mg/L dimethoate 93.17mg/L dimethoate respectively. $n=8$ for both groups.

3.4 Discussion

3.4.1 AChE activity – a marker of pesticide exposure

The AChE activity was detected in the three tissues tested in rainbow trout, with the activity levels ranked as: muscle AChE > brain AChE > plasma AChE. There was no significant activity of BChE, as previously shown for steelhead trout (*Oncorhynchus mykiss*, Sandahl & Jenkins 2002) and Coho Salmon (*Oncorhynchus kisutch*, Sandahl et al. 2005). Exposure to Dimethoate for 21 days inhibited AChE activity in all three tissues; muscle AChE activity was inhibited the most while brain AChE activity was inhibited the least. Chemical-specific patterns of AChE inhibition may exist: Abbas and Hayton (1997) reported that rainbow trout brain AChE activity was more sensitive than plasma AChE activity to another OP pesticide Paraoxon; however they did not measure muscle AChE. Muscle AChE activity may be more sensitive than brain AChE activity to dimethoate exposure as it takes longer for muscle AChE to recover from pesticide exposures compared to brain (Carr et al. 1997). Our data indicate that inhibition of plasma AChE activity, an endpoint that can be measured with non lethal sampling, is a sensitive marker of exposure to Dimethoate. Muscle AChE was a more sensitive marker, compared to brain AChE, both obtained from lethal sampling. Further research is needed to determine the mechanisms of tissue-specific patterns of AChE inhibition and which type of AChE inhibition causes greatest damage to fish health. In estuarine fish, brain AChE inhibition > 70% was associated with mortality, and sublethal effects on stamina were detected with inhibitions as low as 50% (Fulton & Key 2001). In our study with rainbow trout, inhibition of brain AChE as high as 87.91% was detected in the exposure to Dimethoate at 0.1% LC50, with no dimethoate-exposure dependent effect on condition

factor, liver somatic index or mortality. These results suggest that even though AChE was greatly impacted, fish survived and were still feeding normally. Major differences in tolerance to AChE inhibition seem to exist between fish, and although these differences are important and may determine which species survive pesticide exposures in the field, our understanding of mechanisms mediating these differences remains limited at present.

Temperature was not a factor in AChE inhibition in our study with rainbow trout; however, the warming regime used may not have challenged the fish sufficiently. The effects of temperature on AChE inhibition vary with species and measurement conditions. It has been suggested that to cause a significant change in AChE activity in rainbow trout, water temperature needs to change by at least 12°C, with the lower temperature below 10°C (Beauvais et al. 2002). In our study, the temperature regimes were selected to mimic natural water fluctuations in local rivers (Chapter 2). There was a 10°C difference between warm and cold water treatment groups for the first 10 days, with daily fluctuations of 5-6°C over the cold water group for the next 11 days. The scale and temporal pattern of temperature fluctuations in our study may have been too small to affect AChE activity. Future studies will further investigate the effects of temperature fluctuations on AChE activity in rainbow trout.

3.4.2 Physiological Stress Response

Exposure to Dimethoate decreased plasma cortisol levels in rainbow trout, suggesting that dimethoate disrupted the hypothalamo-pituitary-interrenal pathways leading to production of cortisol. In a mammalian study, Walsh et al. (2000) demonstrated that Dimethoate blocked the steroidogenic pathway in the testis of rats by inhibiting the StAR protein responsible for shuttling cholesterol across the mitochondria

and making it available for steroid synthesis. The decreased plasma cortisol in trout exposed to Dimethoate in the present study is consistent with the effects of Dimethoate in rat testis. However, when the secretory capacity of head kidneys isolated from fish exposed to Dimethoate was tested *in vitro* and the head kidney cells were stimulated with exogenous ACTH, normal cortisol secretion was detected. If the mechanism of cortisol disruption (low plasma cortisol) by Dimethoate involved inhibition of StAR protein, stimulation with ACTH would not cause cortisol secretion as the StAR protein acts downstream from the ACTH step in the steroidogenic pathway. Our *in vitro* results with head kidneys of fish exposed to Dimethoate (pure grade) indicate that the secretory capacity of the interrenal tissue was normal since providing ACTH to the head kidney cells did result in normal secretion of cortisol. To determine whether the low plasma cortisol levels in fish exposed to Dimethoate were caused by disruption of pituitary ACTH, or alterations to cortisol metabolism and half-life, as has been suggested for other pollutants (Vijayan et al. 1997), further investigations would be required. Species differences in mechanisms of Dimethoate endocrine toxicity could be responsible for the difference between the results of Walsh et al. (2000) in rat and our results with fish, however the steroidogenic pathways of different vertebrates are very similar (Vijayan et al. 2005). Differences in chemicals used seem more likely to cause different responses. Walsh et al. (2000) used commercial dimethoate (Cygon) while the current study used pure grade Dimethoate.

To test the hypothesis that differences in Dimethoate formulations, rather than species differences, are responsible for the different effects on steroidogenesis, an *in vitro* approach was used in the present study to investigate the effects of two types of

Dimethoate, pure grade and the commercial form of Dimethoate, Cygon, on cortisol secretion in rainbow trout. The pure grade Dimethoate, same chemical as the one used in the *in vivo* experiment, had no effect on cortisol secretion or cell viability when head kidney cells isolated from uncontaminated trout were exposed *in vitro*. The results confirmed that pure grade Dimethoate was not blocking cortisol secretion or affecting StAR protein. The LC₅₀ *in vitro* (concentration that kills 50% of cells) and the EC₅₀ *in vitro* (concentration that decreases cortisol by 50%) could not be determined. In contrast, *in vitro* exposure to Cygon, caused a decrease in cortisol secretion and a reduction in cell viability. Commercial grade Dimethoate has the active ingredient of pure grade Dimethoate but also contains proprietary (unspecified) ingredients such as surfactants. Some studies have shown that commercial forms of pesticides, with surfactants added, are more toxic than the pure products (Beusen & Neven 1989), either by crossing the cells membranes faster or by facilitating transformation into toxic metabolites. Many OPs need to be biotransformed into an oxon form by the liver before they can exert their toxic effects (Grue et al. 1991, Hoffman & Papendorf 2006). Pure grade Dimethoate did not have an effect on cell viability or cortisol secretion *in vitro* because it may require transformation by the liver or it was not able to cross the cell membranes.

The addition of the proprietary ingredients rendered Dimethoate, in the form of Cygon, toxic to the head kidney cells and cortisol production. The EC₅₀, concentration of Cygon at which 50% of cortisol secretion was impaired, at 11°C was lower than the LC₅₀, indicating that the concentration required to disrupt cortisol synthesis was lower than the concentration that killed cells. Thus at 11°C, the secretory processes, possibly function of the StAR protein, were more sensitive to the toxicity of

Cygon than the processes controlling cell viability. Our study also showed that the EC_{50} at 11°C was lower than the EC_{50} at 20°C, providing evidence that the secretory processes were more vulnerable to Cygon at the colder temperature. However at 20°C the EC_{50} was not different from the LC_{50} , suggesting that cortisol secretion was decreased because of mortality of the secretory cells, rather than functional disruption of the secretory processes. Further experimentation is needed to identify the intracellular steps targeted by Dimethoate within the signalling pathway leading to cortisol. In vitro experiments using pure grade or the commercial Dimethoate demonstrated that even though there were no differences in cell viability at 11°C and 20°C, cortisol secretions were consistently higher at 20°C. These results suggest that the vulnerability of cortisol secretion to the pesticide is higher at the colder temperature, even though rainbow trout is a cold water fish species. A previous study by Leblond et al. (2001) reported that cortisol secretion in rainbow trout head kidney cells was lower at 25 and 35°C compared to 5 or 15°C. Further experimentation using a wide range of temperatures is needed to determine the impact that temperature has on cortisol secretion and the stress response.

The physiological stress response is characterized not only by increased plasma cortisol but also by increased plasma glucose and decreased liver glycogen. Dimethoate did not have an effect on plasma glucose or liver glycogen, suggesting that either Dimethoate was not acting as a chemical stressor, or the pathway in the stress response was blocked. These results are consistent with another study where there were no changes in plasma glucose or muscle glycogen in Nile tilapia (*Oreochromis niloticus* L.) exposed to Dimethoate (Sweilum 2006). In our study with rainbow trout, Na^+/K^+ ATPase activity increased with increasing Dimethoate exposure, suggesting that

Dimethoate may alter osmoregulation. Na^+/K^+ ATPase is the enzyme responsible for maintaining transmembrane gradients of Na^+ and K^+ , and changes in the enzyme activity provide a measurement of ion-balance disturbances and the capacity of the fish to osmoregulate under chemical stress (Frasco & Guilhermino 2002). Our results are first to report a change in this enzyme in rainbow trout exposed to Dimethoate. A previous study with *Poecilia reticulata* reported no change in Na^+/K^+ ATPase activity in fish exposed to Dimethoate (Frasco & Guilhermino 2002).

Temperature fluctuations used in this study did not have an effect on plasma cortisol and glucose, or liver glycogen. Water temperature outside the thermal preference of a fish has been reported to alter the stress response (Brett 1971, Bartell et al. 1968, Brandt 1993). Warm temperatures increased plasma cortisol (Strange 1980, Davis et al. 1984, Davis & Parker 1990, Davis 2004) and glucose (Carmichael 1984, Van Dijk et al. 1999, Lerman 2004), and decreased liver glycogen (Viant et al. 2003, Lerman 2004). However, a decrease in plasma glucose at warm water temperatures has been also observed (Davis 2004). Carmichael (1984) found that a change of 12°C (from 10 to 22°C) caused an increase in plasma cortisol and glucose, while a change of 6°C (from 10 to 16°C) caused no change. The change in temperature in our study may not have been high enough or sustained long enough to cause a detectable change in the stress responses.

3.5 Conclusions

Exposure to Dimethoate, both *in vivo* and *in vitro*, had significant effects in rainbow trout in the present study. The activity of AChE was inhibited in muscle, plasma, and

brain and while fish did survive a 21 day exposure to 5% LC₅₀ of dimethoate, AChE was inhibited as much as 90%. Such inhibition would ultimately, upon longer exposures, cause behavioural effects. Dimethoate disrupted the physiological stress response, causing low plasma cortisol levels in vivo, however further studies are required to determine the exact mechanism of the inhibition. Pure grade Dimethoate, *in vitro*, did not cause a decrease in viability and cortisol secretion while commercial Dimethoate disrupted cortisol secretion and decreased viability. Further toxicity testing of Dimethoate should focus on commercial Dimethoate, the chemical applied in the field. Further investigations are required to determine what causes commercial Dimethoate to be more toxic than pure grade Dimethoate.

3.6 References

- Abbas, R. & Hayton, W.L. 1997. A physiologically based pharmacokinetic and pharmacodynamic model for paraoxon in rainbow trout. *Toxicology and Applied Pharmacology* 145:192-201.
- Athanasopoulos, P., Kyriakidis, N.B., & Georgitsanakou, I. 2000. Effect of storage temperature on the degradation of dimethoate in fortified orange and peach juices. *Journal of Agriculture and Food Chemistry* 48:4896-4899.
- Bartell, S.M., Breck, J.E., Gardner, R.H., & Brenkert, A.L. 1986. Individual parameter perturbation and error analysis of fish bioenergetics models. *Canadian Journal of Fisheries and Aquatic Science* 43:160-168.
- Barton, B.A. & Peter, R.E. 1982. Plasma cortisol stress response in fingerling rainbow trout *Salmo gairdneri* Richardson, to various transport conditions, anaesthesia, and cold shock. *Journal of Fish Biology* 20:39-51.
- Beauvais, S.L., Cole, K.J., Atchison, G.J., & Coffey, M. 2002. Factors affecting brain cholinesterase activity in bluegill (*Lepomis macrochirus*). *Water, Air, and Soil Pollution* 135:249-264.
- Beauvais, S.L., Jones, S.B., Parris, J.T., Brewer, S.K., & Little, E.E. 2001. Cholinergic and behavioural neurotoxicity of carbaryl and cadmium to larval rainbow trout (*Oncorhynchus mykiss*). *Ecotoxicology and Environmental Safety* 49:84-90.
- Begum, G. & Vijayaraghavan, S. 1995. Chronic effects of dimethoate on the reproductive potential of the fresh-water teleost, *Clarias batrachus*. *Pesticide Science* 44:233-236.
- Begum, G. & Vijayaraghavan, S. 1999. Effect of acute exposure of the organophosphate insecticide Rogor on some biochemical aspects of *Clarias batrachus* (Linnaeus). *Environmental Research* 80A:80-83.
- Beusen, J.-M. & Neven, B. 1989. Toxicity of dimethoate to *Daphnia magna* and freshwater fish. *Bulletin of Environmental Contamination and Toxicology* 42:126-133.
- Bhattacharya, S. 1993. Target and non-target effects of anticholinesterase pesticides in fish in: *Proceeding of the second European conference on toxicology. Science of the Total Environment Supplement*:859-866.
- Bleau, H., Daniel, C., Chevalier, G., van Tra, H., & Hontela, A. 1996. Effects of acute exposure to mercury chloride and methylmercury on plasma cortisol, T3, T4,

- glucose, and liver glycogen in rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology* 34:221-245.
- Bloomfield, J.P., Williams, R.J., Gooddy, D.C., Cape, J.N., & Guha, P. 2006. Impacts of climate change on the fate and behaviour of pesticides in surface and groundwater – a UK perspective. *Science of the total Environment* 369:163-177.
- Brandt, S.B. 1993. The effect of thermal fronts on fish growth: A bioenergetics evaluation of food and temperature. *Estuaries* 16:142-159.
- Brett, J.R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). *American Zoologist* 11:99-113.
- Carmichael, G.J., Tomasso, J.R., Simco, B.A., & Davis, K.B. 1984. Confinement and water quality-induced stress in largemouth bass. *Transactions of the American Fisheries Society* 113:767-777.
- Carr, R.L., Ho, L.L., & Chambers, J.E. 1997. Selective toxicity of chlorpyrifos to several species of fish during an environmental exposure: Biochemical mechanisms. *Environmental Toxicology and Chemistry* 16: 2369-2374.
- Chuiko, G.M. 2000. Comparative study of acetylcholinesterase and butyrylcholinesterase in brain and serum of several freshwater fish: specific activities and in vitro inhibition by DDVP, an organophosphorus pesticide. *Comparative Biochemistry and Physiology* 127C:233-242.
- Davis, K.B. 2004. Temperature affects physiological stress responses to acute confinement in sunshine bass (*Morone chrysops* X *Morone saxatilis*). *Comparative Biochemistry and Physiology* 139A:433-440.
- Davis, K.B. & Parker, N.C. 1990. Physiological stress in striped bass: effect of acclimation temperature. *Aquaculture* 91:349-358.
- Davis, K.B. & Peterson, B.C. 2006. The effect of temperature, stress, and cortisol on plasma IFG-I and IGFBPs in sunshine bass. *General and Comparative Endocrinology* 149:219-225.
- Davis, K.B., Stuttle, M.A., & Parker, N.C. 1984 Biotic and abiotic influences on corticosteroid hormone rhythms in channel catfish. *Transactions of the American Fisheries Society* 118:414-421.
- Dell’Omo, G., Bryenton, R., & Shore, R.F. 1997. Effects of exposure to an organophosphate pesticide on behaviour and acetylcholinesterase activity in the common shrew, *Sorex araneus*. *Environmental Toxicology and Chemistry* 16:272-276.

- De Mel, G.W.J.L.M.V.T.M. & Pathiratne, A. 2005. Toxicity assessment of insecticides commonly used in rice pest management to the fry of common carp, *Cyprinus carpio*, a food fish culturable in rice fields. *Journal of Applied Ichthyology* 21:146-150.
- De Silva, P.M.C.S. & Samayawardhena, L.A. 2002. Low concentrations of Lorsban in water result in far reaching behavioral and histological effects in early life stages in guppy. *Ecotoxicology and Environmental Safety* 53:248-254.
- Dorval, J., Leblond, V., Deblois, C., & Hontela, A. 2005. Oxidative stress and endocrine endpoints in white sucker (*Catostomus commersoni*) from a river impacted by agricultural chemicals. *Environmental Toxicology and Chemistry* 24:1273-1280.
- Ellman, G.L., Courtney, K.D., Anders, V., Jr., & Featherstone, R.M. 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 7:88-95.
- Farag, A.T., Karkour, T. A.-Z., & Okazy, A. 2006. Developmental toxicity of orally administered technical dimethoate in rats. *Birth Defects Research* 77B:40-46.
- Frasco, M.F. & Guilhermino, L. 2002. Effects of dimethoate and beta-naphthoflavone on selected biomarkers of *Poecilia reticulata*. *Fish Physiology and Biochemistry* 26:149-156.
- Fulton, M.H. & Key, P.B. 2001. Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticide exposure and effects. *Environmental Toxicology and Chemistry* 20:37-45.
- Gill, T.S., Pant, J.C., & Pant, J. 1988. Gill, liver, and kidney lesions associated with experimental exposures to carbaryl and dimethoate in the fish (*Puntius conchonius* Ham.). *Bulletin of Environmental Contamination and Toxicology* 41:71-78.
- Grue, C.E., Hart, A.D.M., & Mineau, P. 1991. Biological consequences of depressed brain cholinesterase activity in wildlife In: *Cholinesterase-inhibiting insecticides* ed. Mineau, P. Elsevier. Amsterdam.
- Hoffmann, U. & Papendorf, T. 2006. Organophosphate poisonings with parathion and dimethoate. *Intensive Care Medicine* 32:464-486.
- Hogan, J.W. 1970. Water temperature as a source of variation in specific activity of brain acetylcholinesterase of bluegills. *Bulletin of Environmental Contamination & Toxicology* 5:347-353.
- Holliday, C.W. 1985. Salinity-induced changes in gill Na,K-ATPase activity in the mud fiddler crab, *Uca pugnax*. *Journal of Experimental Zoology* 233:199-208.

- Hontela, A. 1998. Interrenal dysfunction in fish from contaminated sites: In vivo and in vitro assessment. *Environmental Toxicology and Chemistry* 17:44-48.
- Johnson, W.W. & Finley, M.T. 1980. Handbook of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates. Resource Publication 137. U.S. Department of the Interior, Fish and Wildlife Service, Washington, DC, 4-17
- Leblond, V.S., Bisson, M. & Hontela, A. 2001. Inhibition of cortisol secretion in dispersed head kidney cells of rainbow trout (*Oncorhynchus mykiss*) by endosulfan, an organochlorine pesticide. *General and Comparative Endocrinology* 121:48-56.
- Lermen, C.L., Lappe, R., Crestani, M., Vieira, V.P., Gioda, C.R., Schetinger, M.R.C., Baldissotto, B., Moraes, G., & Morsch, V.M. 2004. Effect of different temperature regimes on metabolic and blood parameters of silver catfish *Rhamdia quelen*. *Aquaculture* 239:497-507.
- Levesque, H.M., Dorval, J., Van Der Kraak, G.J., Campbell, P.G.C. & Hontela, A. 2003. Hormonal, morphological, and physiological responses of yellow perch (*Perca flavescens*) to chronic environmental metal exposures. *Journal of Toxicology and Environmental Health* 66A:657-676.
- Little, E.E. & Finger, S.E. 1990. Swimming behavior as an indicator of sublethal toxicity in fish. *Environmental Toxicology and Chemistry* 9:13-19
- Martikainen, E. & Rantalainen, M.-L. 1999. Temperature-time relationship in collembolan response to chemical exposure. *Ecotoxicology and Environmental Safety* 42:236-244.
- Morgan, I.J., D'Cruz, L.M., Dockray, J.J., Linton, T.K., McDonald, D.G., & Wood, C.M. 1998. The effect of elevated winter temperature and sub-lethal pollutants (low pH, elevated ammonia) on protein turnover in the gill and liver of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiology and Biochemistry* 19:377-389.
- Phillips, T.A., Summerfelt, R.C., & Atchison, G.J. 2002. Environmental, biological, and methodological factors affecting cholinesterase activity in walleye (*Stizostedion vitreum*). *Archives of Environmental Contamination and Toxicology* 43:75-80.
- Sandahl, J.F., Baldwin, D.H., Jenkins, J.J., & Scholz, N.L. 2005. Comparative thresholds for acetylcholinesterase inhibition and behavioral impairment in coho salmon exposed to chlorpyrifos. *Environmental Toxicology and Chemistry* 24:136-145
- Sandahl, J.F. & Jenkins, J.J. 2002. Pacific steelhead (*Oncorhynchus mykiss*) exposed to chlorpyrifos: Benchmark concentration estimates for acetylcholinesterase inhibition. *Environmental Toxicology and Chemistry* 21:2452-2458.

- Schreck, C.B., Contreras-Sanchez, W., & Fitzpatrick, M.S. 2001. Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture* 197:3-24.
- Strange, R.J. 1980. Acclimation temperature influences cortisol and glucose concentrations in stressed channel catfish. *Transactions of the American Fisheries Society* 109:289-303.
- Sweilum, M.A. 2006. Effect of sublethal toxicity of some pesticides on growth parameters, haematological properties and total production of Nile tilapia (*Oreochromis niloticus* L.) and water quality of ponds. *Aquaculture Research* 37:1079-1089.
- Teles, M., Pacheco, M., & Santos, M.A. 2007. Endocrine and metabolic responses of *Anguilla Anguilla* L. caged in a freshwater-wetland (Pateira de Fermentelos-Portugal). *Science of the Total Environment* 372:562-570.
- Thangavel, P., Sumathiral, K., Karthikeyan, S., & Ramaswamy, M. 2005. Endocrine response of the freshwater teleost, *Sarotherodon mossambicus* (Peters) to dimecron exposure. *Chemosphere* 61:1083-1092.
- Tuduri, L. Harner, T., Blanchard, P., Li, Y.-F., Poissant, L., Waite, D.T., Murphy, C., & Belzer, W. 2006. A review of currently used pesticides (CUPs) in Canadian air and precipitation: Part 2: Regional information and perspectives. *Atmospheric Environment* 40: 1563-1578.
- Van Dijk, P.L.M., Tesch, C., Hardewig, I., & Pörtner, H.O. 1999. Physiological disturbances at critically high temperatures: a comparison between stenothermal antarctic and eurythermal temperate eelpouts (*Zoaridae*). *The Journal of Experimental Biology* 202:3611-3621.
- Viant, M.R., Werner, I., Rosenblum, E.S., Gantner, A.S., Tjeerdema, R.S., & Johnson, M.L. 2003. Correlation between heat-shock protein induction and reduced metabolic condition in juvenile steelhead trout (*Oncorhynchus mykiss*) chronically exposed to elevated temperature. *Fish Physiology and Biochemistry* 29:159-171.
- Vijayan M.M., Feist, G., Otto, D.M.E., Schreck, C.B., & Moon, T.W. 1997. 3,3', 4, 4'-tetrachlorobiphenyl affects cortisol dynamics and hepatic function in rainbow trout. *Aquatic Toxicology* 37: 87-98.
- Vijayan, M.M., Prunet, P., & Boone, A.N. 2005. Xenobiotic impact on corticosteroid signaling In: *Biochemistry and Molecular Biology of Fishes Vol 6*. Mommsen, T.P. & Moon, T.W. Eds. Elsevier B.V.

- Walsh, L.P., Webster, D.R., & Stocco, D.M. 2000. Dimethoate inhibits steroidogenesis by disruption transcription of the steroidogenic acute regulatory (StAR) gene. *Journal of Endocrinology* 167:253-263.
- Wauchope, R.D., Buttler, T.M., Hornsby, A.G., Augustijn-Beckers, P.W.M., & Burt, J.P. 1992. The SCS/ARS/CES pesticide properties database for environmental decision-making. *Reviews of environmental contamination and toxicity*. 123:1-155.
- Wendelaar Bonga, S.E. 1997. The stress response in fish. *Physiological Reviews* 77:591-625.
- World Health Organization (WHO). 1989. International program on chemical safety. Environmental health criteria 90: Dimethoate. EHC 90
- Zinkl, J.G., Shea, P.J., Nakamoto, R.J., & Callman, J. 1987. Technical and biological considerations for the analysis of brain cholinesterase of rainbow trout. *Transactions of the American Fisheries Society* 116:570-573.

Chapter 4.

General Discussion and Conclusion

4.1 General Overview

The purpose of this project was to investigate the effects of multiple stressors on the physiological status of fish and determine if the physiological stress response was activated or impaired in response to pesticide exposure and increasing water temperature. Furthermore, the responses of different fish species were compared to better understand the processes that make fish vulnerable to environmental stressors. Field sampling of wild fish (whitefish and suckers) and controlled laboratory exposures of rainbow trout (a well validated fish model) were used in this study that provided new data on physiological markers, including plasma cortisol, liver glycogen, plasma glucose, condition factor, Na⁺/K⁺ ATPase activity and AChE activity in fish.

4.2 Oldman Study Conclusions (Chapter 2)

The objective of the Oldman study was to determine how multiple stressors (pesticides and temperature) affect the physiological status and stress response in cold-water fish (Mountain Whitefish, *Prosopium williamsoni*) and cool-water fish (Suckers, *Catostomus* sp.). The findings of this study determined that the Oldman River has a west-to-east warm temperature gradient and a gradient for 2,4-D and MCP (Mecoprop). However, a gradient was not detected for organophosphates (OPs) or carbamates (CBs), possibly because of the limited testing procedures and a short half-life of these pesticides (Morrison & Wells 1981, Fulton & Key 2001). Although we were unable to detect OPs and CBs in the water, site differences and species differences in the activity of AChE, a marker of exposure to OP and CB were detected. Moreover, there was a significant

difference in AChE activity between whitefish and suckers. Whitefish had lower levels of AChE activity, suggesting either there is a physiological difference between the two species or whitefish had a greater exposure to pesticides (possibly through diet). It is difficult to conclude at this time whether the AChE activity is normal for whitefish and suckers since ours are the first data reported for these species in this system. Further studies using whitefish and suckers are needed to determine how pesticide exposure and temperature affect AChE activity but our results suggest that whitefish may be highly vulnerable to pesticides.

Mountain white fish at sites outside of their thermal preference exhibited a higher stress response, as measured by plasma cortisol levels and liver glycogen compared to suckers at the same sites. The increase in plasma cortisol and the decrease in liver glycogen follow both the temperature and pesticide gradient suggesting one or both of these factors act as a stressor for both whitefish and sucker. Again this was the first study of the physiological stress response in whitefish so it is difficult to determine if the values found are within the normal range. Overall this study found species specific differences between whitefish and suckers in regards to AChE activity, plasma cortisol and liver glycogen. The results suggest that whitefish may be more sensitive to water temperature and pesticide exposure and may be more at risk if water temperature and pesticide usage continue to increase. This study suggests that monitoring of whitefish should continue as there is little information regarding their ability to handle stressors. Further information would help determine if whitefish are at risk and whether management strategies are needed to maintain this population.

4.3 Dimethoate Study Conclusions (Chapter 3)

The objective of the Dimethoate study was to investigate the effects of Dimethoate and temperature on the physiological stress response and AChE activity in rainbow trout (*Oncorhynchus mykiss*). This study also looked at the differences between pure grade Dimethoate and commercial dimethoate (Cygon). In the *in vivo* study, rainbow trout were exposed to low amounts (0.5 and 1% LC₅₀) of Dimethoate; these low concentrations decreased AChE activity in a dose dependent manner in the muscle, brain, and plasma. However, temperature did not have an effect on AChE activity. It has been suggested that in order for changes in AChE activity to occur due to temperature, the change in temperature needs to be at least 10°C and the lower temperature below 12°C (Beauvais et al. 2002). This study could be repeated with a lower starting temperature and a greater change in temperature to determine if temperature is a factor in AChE activity.

Plasma cortisol levels decreased with increasing Dimethoate exposure suggesting that Dimethoate disrupts the physiological stress response production of cortisol. However, when the secretory capacity of the isolated head kidney stimulated with adrenocorticotrophic hormone (ACTH) was assessed, cortisol was produced. This suggests that Dimethoate blocks cortisol production within the hypothalamo-pituitary-interrenal axis at a step before ACTH acts. These results are different from other studies that suggest that Dimethoate disrupts cortisol secretion after ACTH, by blocking the StAR protein (Walsh et al. 2000).

To further determine where Dimethoate blocks cortisol secretion, an *in vitro* system was used. First, the head kidney cells of rainbow trout were exposed to pure grade dimethoate (same as the *in vivo* experiment). Pure grade Dimethoate had no effect on cortisol secretion even at the highest concentration. Cygon, a commercial Dimethoate was then used, similar to the original Walsh et al. (2000) experiment. Commercial dimethoate caused a reduction in both cortisol secretion and cell viability. Cortisol secretion decreased more than cell viability indicating that commercial dimethoate blocks cortisol production at lower concentrations of the pesticide than those killing the cells. This study also demonstrated that there are implications to using pure versus commercial grade products in toxicity testing. Commercial products often have proprietary ingredients that can cause cellular damage or allow the toxicant to infiltrate the cell.

Temperature did have an effect on cortisol secretion *in Vitro* both with pure and commercial grade Dimethoate. The higher temperature increased cortisol secretion however temperature did not have an effect on viability. Temperature also did not have an effect on cortisol secretion in the *in Vivo* experiment. Temperature did not affect plasma glucose or liver glycogen, suggesting that temperature does not affect the stress response or that the temperature change was not great enough to cause a change in the stress response.

4.4 Concluding Statements

This study provided novel information regarding the physiological stress response and acetylcholinesterase. This study also addressed important aspects of multiple stressors affecting fish in southern Alberta and Canada as a whole. Climate change is a

very important issue impacting agriculture, industry and government. Changes in climate will be an added stressor to natural and anthropogenic stressors that are already present in aquatic systems. This thesis attempted to discover how temperature would influence pesticide exposure in freshwater fish and it provided new information how these two stressors affect freshwater fish.

4.5 References

Beauvais, S.L., Cole, K.J., Atchison, G.J., & Coffey, M. 2002. Factors affecting brain cholinesterase activity in bluegill (*Lepomis macrochirus*). *Water, Air, and Soil Pollution* 135:249-264.

Fulton, M.H. & Key, P.B. 2001. Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticide exposure and effects. *Environmental Toxicology and Chemistry* 20:37-45.

Morrison, B.R.S. & Wells, D.E. 1981. The fate of fenitrothion in a stream environment and its effect on the fauna, following aerial spraying of a Scottish forest. *The Science of the Total Environment* 19:233-252.

Walsh, L.P., Webster, D.R., & Stocco, D.M. 2000. Dimethoate inhibits steroidogenesis by disruption transcription of the steroidogenic acute regulatory (StAR) gene. *Journal of Endocrinology* 167:253-263.

Appendix

Table A.1. AChE activity in Salmonids and Catostomids exposed to organophosphates (OP), carbamates (CB), or field pesticides

| Pesticide | Exposure | Dose | Specific Activity | Species | Tissue | Source |
|-----------|-----------------|-------------|--------------------------------|-------------------------------|--------|-----------------------|
| | in water | | 9.28 AChE (mmol/min/g tissue) | <i>O. mykiss</i> ^a | Brain | Davies et al. 1994 |
| | in water | | 192 (nmol/min/mg protein) | <i>O. mykiss</i> | Brain | Shao-nan&De-fang 1996 |
| | <i>in Vitro</i> | | 225 AChE (nmol/min/mg protein) | <i>O. mykiss</i> | Brain | Shaonan et al. 2004 |
| Controls | in water | No exposure | 16.06 AChE (nmol/g wet tissue) | <i>O. mykiss</i> | Brain | Szabó et al. 1991 |
| | in water | | 4.66 AChE (nmol/g wet tissue) | <i>O. mykiss</i> | Gut | |
| | in water | | 1.47 AChE (nmol/g wet tissue) | <i>O. mykiss</i> | Heart | |
| | in water | | 7.08 AChE (mmol/min/g tissue) | <i>O. mykiss</i> | Muscle | Davies et al.1994 |
| | in water | | 3.32 AChE (nmol/g wet tissue) | <i>O. mykiss</i> | Muscle | Szabó et al. 1991 |

^aRainbow trout, *Oncorhynchus mykiss*

| Pesticide | Exposure | Dose | Effect | Species | Tissue | Source |
|----------------------|----------|--------------|---|------------------------------------|------------------|----------------------|
| Aldicarb (CB) | in water | 0.5 mg/L | 75% AChE inhibition | <i>O. mykiss</i> | Brain | Wang et al. 2001 |
| | in water | 0.5 mg/L | 30% AChE inhibition | <i>O. mykiss</i> | Muscle | |
| Azamethiphos (OP) | in water | 0.1 ppm | 83% AChE inhibition (60 min exposure) | <i>O. mykiss</i> | Brain | Intorre et al. 2004 |
| | in water | 0.1 ppm | 79 % AChE inhibition (120 min exposure) | <i>O. mykiss</i> | Brain | |
| | in water | 0.1 ppm | 62% AChE inhibition (240 min exposure) | <i>O. mykiss</i> | Brain | |
| Azinphos methyl (OP) | in water | 0.001 mg/L | 70% AChE inhibition | <i>O. mykiss</i> | Brain | Ferrari et al. 2004 |
| Carbaryl (CB) | in water | 188-750 µg/L | 14-38% AChE inhibition for 24 hr exposure | <i>O. mykiss</i> | Brain | Beauvais et al. 2001 |
| | in water | 188-750 µg/L | 32-41% AChE inhibition for 96 hr exposure | <i>O. mykiss</i> | Brain | |
| | in Vivo | 188-750 µg/L | 7-14% AChE inhibition for 48 hr recovery | <i>O. mykiss</i> | Brain | |
| | in Vivo | 0.5 mg/L | 77% AChE inhibition | <i>O. mykiss</i> | Brain | Ferrari et al. 2004 |
| | in Vivo | 66 µg/L | 50% AChE inhibition | <i>O. tshawytscha</i> ^b | Olfactory tissue | Scholz et al. 2006 |

| Pesticide | Exposure | Dose | Effect | Species | Tissue | Source |
|----------------------|-----------------|-----------------|---|--------------------------------|-------------------|------------------------|
| Carbofuran (CB) | <i>in Vitro</i> | 14.7 µg/L | 32.4% AChE inhibition | <i>O. kisutch</i> ^c | Olfactory rosette | Jarrard et al. 2004 |
| | in water | 200 µg/L | 50% AChE inhibition | <i>O. tshawytscha</i> | Olfactory tissue | Scholz et al. 2006 |
| Chlorpyrifos (OP) | in water | 0.6 µg/L | 23.1% AChE inhibition | <i>O. kisutch</i> | Brain | Sandahl et al. 2005 |
| | in water | 2.5 µg/L | 64% AChE inhibition | <i>O. kisutch</i> | Brain | |
| | in water | 0.6 µg/L | 12% AChE inhibition | <i>O. kisutch</i> | Muscle | |
| | in water | 2.5 µg/L | 67% AChE inhibition | <i>O. kisutch</i> | Muscle | |
| | in water | 0.625-2.50 µg/L | 7-38.4% AChE inhibition | <i>O. mykiss</i> | Brain | Sandahl & Jenkins 2002 |
| | in water | 1.2 µg/L | 8% AChE inhibition | <i>O. tshawytscha</i> | Brain | Eder et al. 2004 |
| | in water | 7.3 µg/L | 15% AChE inhibition | <i>O. tshawytscha</i> | Brain | |
| | in water | 7.3 µg/L | 85% AChE inhibition compared to solvent control | <i>O. tshawytscha</i> | Brain | Wheelock et al. 2005 |
| | in water | 7.3 µg/L | 8% AChE inhibition | <i>O. tshawytscha</i> | Muscle | Eder et al. 2004 |
| | in water | 7.3 µg/L | 93% AChE inhibition compared to solvent control | <i>O. tshawytscha</i> | Muscle | Wheelock et al. 2005 |

| Pesticide | Exposure | Dose | Effect | Species | Tissue | Source |
|------------------------|----------|---------------|---|--------------------------------|------------------|--------------------------|
| Chlorpyrifos-oxon (OP) | in water | 8 µg/L | 50% AChE inhibition | <i>O. tshawytscha</i> | Olfactory tissue | Scholz et al. 2006 |
| | in water | 5 µg/L | 50% AChE inhibition | <i>Catostomus occidentalis</i> | Brain | Whitehead et al. 2005 |
| Diazinon (OP) | in water | 4 µg/L | 50% AChE inhibition | <i>Catostomus occidentalis</i> | Muscle | |
| | in water | 250-1000 µg/L | 16-37% AChE inhibition | <i>O. mykiss</i> | Brain | Beauvais et al. 2000 |
| | in water | 2.5 µM | 50% AChE inhibition | <i>O. mykiss</i> | Brain | Keizer et al. 1995 |
| Diazinon-oxon (OP) | in water | 525 µg/L | 50% AChE inhibition | <i>O. tshawytscha</i> | Olfactory tissue | Scholz et al. 2006 |
| Dichlorvos (OP) | in water | 1.136 ppm | 45-58% AChE inhibition (depending on oxygen levels) | <i>O. mykiss</i> | Brain | Høy et al. 1991 |
| | in water | 1300 µg/L | 63% AChE inhibition 7 day exposure | <i>Cyprinus carpio*</i> | Brain | De Mel & Pathiratne 2005 |
| Dimethoate (OP) | in water | 1301 µg/L | 69% AChE inhibition 14 day exposure | <i>Cyprinus carpio*</i> | Brain | |
| | in water | 1 mg/L | 70% AChE inhibition | <i>Poecilia reticulata*</i> | Head | Frasco & Guihermino 2002 |
| | in water | 0.125 mg/L | 50% AChE inhibition | <i>Poecilia reticulata*</i> | Head | |

| Pesticide | Exposure | Dose | Effect | Species | Tissue | Source |
|---------------------|-----------------|----------------|---|-----------------------|------------------|------------------------|
| Fenitrothion (OP) | in water | 15 to 220 µg/L | 1.68- 71.9% AChE inhibition for 10 d exposure | <i>O. mykiss</i> | Brain | Davies et al.1994 |
| | in water | 15 to 220 µg/L | 41.2- 83.9% AChE inhibition for 20 d exposure | <i>O. mykiss</i> | Brain | |
| IPBC (CB) | <i>in Vitro</i> | 0.0047 µg/L | 196% AChE activity of control | <i>O. kisutch</i> | Brain | Jarrard et al. 2004 |
| | <i>in Vitro</i> | 0.47 µg/L | 175% AChE activity of control | <i>O. kisutch</i> | Brain | |
| | <i>in Vitro</i> | 47.5 µg/L | 186% AChE activity of control | <i>O. kisutch</i> | Brain | |
| Malathion (OP) | in water | 20-40 µg/L | 19.2-63% AChE inhibition | <i>O. mykiss</i> | Brain | Beauvais et al. 2000 |
| Malathion-oxon (OP) | in water | 4.1 µg/L | 50% AChE inhibition | <i>O. tshawytscha</i> | Olfactory tissue | Scholz et al. 2006 |
| Mancozeb (CB) | <i>in Vitro</i> | 0.0022 µg/L | 187% AChE activity of control | <i>O. kisutch</i> | Brain | Jarrard et al. 2004 |
| | <i>in Vitro</i> | 0.022 µg/L | 143% AChE activity of control | <i>O. kisutch</i> | Brain | |
| Methidathion (OP) | in water | 1 µg/L | 22.2% AChE inhibition | <i>O. mykiss</i> | Muscle | Flammarion et al. 1996 |
| | in water | 5 µg/L | 90.5% AChE inhibition | <i>O. mykiss</i> | Muscle | |

| Pesticide | Exposure | Dose | Effect | Species | Tissue | Source |
|----------------------------|-------------|----------|---|------------------------------|--------|----------------------|
| Paraoxon (OP) | in water | 75 ng/ml | 58.9% AChE inhibition after 24 hr | <i>O. mykiss</i> | Brain | Abbas & Hayton 1997 |
| | in water | 76 ng/ml | 51.2% AChE inhibition after 24 hr | <i>O. mykiss</i> | Heart | Abbas & Hayton 1997 |
| | in water | 77 ng/ml | 49.6% AChE inhibition after 24 hr | <i>O. mykiss</i> | Liver | |
| Solvent Control (methanol) | in water | | 11% AChE inhibition | <i>O. tshawytscha</i> | Brain | Wheelock et al. 2005 |
| Urban Rivers | field study | | no inhibition compared to reference site | <i>Salmo trutta</i> | Brain | Payne et al. 1996 |
| | field study | | 40-50% AChE inhibition compared to reference site | <i>Salmo trutta</i> | Muscle | |
| Yamaska River | field study | | 39.1-40.8% AChE inhibition compared to reference site | <i>Catostomus commersoni</i> | Plasma | Dorval et al. 2005 |

^bChinook salmon - *Oncorhynchus tshawytscha*

^cCoho salmon - *Oncorhynchus kisutch*

*data on AChE inhibition in salmonids or catostomids exposed to Dimethoate were not available

| Table A.2. Physiological indicators of stress in fish exposed to pesticides | | | | | | |
|---|-------------|-----------------|------------------|---|----------------------------|--|
| Pesticide | Action | Exposure | Dose | Effect | Species | Source |
| Acephate (OP) | Insecticide | | 0.2-1.3 mg/L | no change in plasma glucose | <i>Galaxias maculatus</i> | Davies et al. 1994 |
| | | in water | 4.4 mg/L | plasma glucose ↑ | <i>Galaxias maculatus</i> | |
| | | in water | 0.2 mg/L | plasma glucose ↑ | <i>Oncorhynchus mykiss</i> | |
| | | in water | 0.4-4.4 mg/L | no change in plasma glucose | <i>Oncorhynchus mykiss</i> | |
| Atrazine (TR) | Herbicide | <i>in Vitro</i> | 500 μM | cortisol secretion ↑ | <i>Oncorhynchus mykiss</i> | Bisson & Hontela 2002 |
| | | <i>in Vitro</i> | 5 to 0.005 μM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | |
| | | in water | 340 μg/L | plasma glucose ↑ | <i>Oncorhynchus mykiss</i> | Davies et al. 1994 Waring & Moore 2004 |
| | | in water | 1.1- 22.7 μ/L | plasma cortisol ↑ | <i>Salmo salar</i> | |
| | | in water | 1-10 μg/L | Na ⁺ -K ⁺ -ATPase ↓ | <i>Salmo salar</i> | |
| Carbofuran (CB) | Insecticide | in water | 50 μg/L | plasma glucose ↑ 150% | <i>Carassius auratus</i> | Bretaud et al. 2002 |
| | | in water | 500 μg/L | plasma glucose ↑ 230% | <i>Carassius auratus</i> | |
| | | in water | 50 μg/L | liver glycogen ↑ 122% | <i>Clarassius auratus</i> | |
| | | in water | 500 μg/L | liver glycogen ↓ 46% | <i>Clarassius auratus</i> | |
| | | in water | 7.66 mg/L | liver glycogen ↓ | <i>Clarias botrachus</i> | Begum 2004 |
| Carbaryl (CB) | Insecticide | in water | 4.0-8.0 mg/L | liver glycogen ↓ | <i>Colisa fasciatus</i> | Singh et al. 2004 |
| | | in water | 4.0-8.0 mg/L | muscle glycogen ↓ | <i>Colisa fasciatus</i> | |
| Pesticide | Action | Exposure | Dose | Effect | Species | Source |

| | | | | | | |
|-------------------|-------------|-----------------|----------------------|--|---------------------------------|-----------------------------|
| Cyanazine (TR) | Herbicide | in water | 900-2000 µg/L | plasma glucose ↓ | <i>Oncorhynchus mykiss</i> | Davies et al. 1994 |
| Cypermethin (PY) | Insecticide | in water | 0.49 µg/L | plasma glucose ↓ | <i>Oncorhynchus mykiss</i> | |
| Deltamethrin (PY) | Insecticide | in water | 0.005 mg/L | liver glycogen ↓ | <i>Clarias gariepinus</i> | Datta & Kaviraj 2003 |
| | | in water | 0.005 mg/L | plasma glucose ↑ | <i>Clarias gariepinus</i> | |
| Diazinon (OP) | Insecticide | <i>in Vitro</i> | 500 µM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Bisson & Hontela 2002 |
| Dimecron (OP) | Insecticide | in water | 0.001 mL/L | serum cortisol ↓ | <i>Saratherodon mossamuicus</i> | Thangavel et al. 2005 |
| | | in water | 21.66 mg/L | muscle glycogen ↓ | <i>Clarias batrachus</i> | Begum & Vijayaraghavan 1999 |
| Dimethoate (OP) | Insecticide | in water | 5-20 mg/L | no change in muscle glycogen | <i>Oreochromis niloticus</i> L. | Sweilum 2006 |
| | | in water | 5-20 mg/L | no change in plasma glucose | <i>Oreochromis niloticus</i> L. | |
| | | in water | 1 mg/L | no change in Na ⁺ /K ⁺ -ATPase | <i>Poecilia reticulata</i> | Frasco & Guihermino 2002 |
| Endosulfan (OC) | Insecticide | <i>in Vitro</i> | 500 µM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Bisson & Hontela 2002 |
| | | <i>in Vitro</i> | 50 µM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | |
| | | <i>in Vitro</i> | 1x10 ⁻⁵ M | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Dorval & Hontela 2003 |
| | | <i>in Vitro</i> | 1x10 ⁻⁵ M | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Dorval et al. 2003 |
| | | <i>in Vitro</i> | 1x10 ⁻⁵ M | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Leblond et al. 2001 |
| | | <i>in Vitro</i> | 1x10 ⁻⁴ M | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | |

| Pesticide | Action | Exposure | Dose | Effect | Species | Source |
|-------------------|-------------|-----------------|--------------|-----------------------------|---------------------------------|-----------------------|
| Fenitrothion (OP) | Insecticide | in water | 0.02 ppm | no change in liver glycogen | <i>Anguilla anguilla</i> | Sancho et al. 1997 |
| | | in water | 0.02 ppm | plasma glucose ↑ | <i>Anguilla anguilla</i> | |
| | | in water | 0.04 ppm | liver glycogen ↓ | <i>Anguilla anguilla</i> | Sancho et al. 1998 |
| | | in water | 0.04 ppm | plasma glucose ↑ | <i>Anguilla anguilla</i> | |
| Folidol 600 (OP) | Insecticide | in water | 2 ppm | liver glycogen ↑ 84% | <i>Brycon cephalus</i> | de Aguiar et al. 2004 |
| | | in water | 2 ppm | glucose ↑ 64% | <i>Brycon cephalus</i> | |
| Malathion (OP) | Insecticide | in water | 1.0-3.0 mg/L | liver glycogen ↓ | <i>Colisa fasciatus</i> | Singh et al. 2004 |
| | | in water | 1.0-3.0 mg/L | muscle glycogen ↓ | <i>Colisa fasciatus</i> | |
| | | in water | 0.5-2.0 mg/L | muscle glycogen ↓ | <i>Oreochromis niloticus L.</i> | Sweilum 2006 |
| | | in water | 0.5-2.0 mg/L | plasma glucose ↑ | <i>Oreochromis niloticus L.</i> | |
| Mancozeb (EB) | Fungicide | <i>in Vitro</i> | 50 μM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Bisson & Hontela 2002 |

| Pesticide | Action | Exposure | Dose | Effect | Species | Source |
|---------------------|-------------|-----------------|----------|--|------------------------------|------------------------|
| Monocrotophos (OP) | Insecticide | in water | 10 mg/L | liver glycogen ↓ 2.4% | <i>Channa punctatus</i> | Samuel & Sastry 1989 |
| | | in water | 10 mg/L | muscle glycogen ↓ 35.4% | <i>Channa punctatus</i> | |
| | | in water | 10 mg/L | plasma glucose ↓ 12.7% | <i>Channa punctatus</i> | |
| <i>o'p</i> DDD (OC) | | <i>in Vitro</i> | 75-200µM | cortisol secretion ↓ | <i>Oncorhynchus mykiss</i> | Lacroix & Hontela 2003 |
| Yamaska River | | Field study | | plasma cortisol ↓ compared to reference site | <i>Catostomus commersoni</i> | Dorval et la. 2005 |
| | | Field study | | liver glycogen ↑ compared to reference site | <i>Catostomus commersoni</i> | |

Pesticide abbreviations: CB - carbamate, EB - ethylene bisdithiocarbamate, OC - organochlorine, OP - organophosphate, PY - pyrethroid, TR – triazine

Table A.3. Behavioral changes in fish exposed to organophosphate (OP) and carbamate (CB) pesticides

| Pesticide | Dose (exposure period) | Affected Behaviour | Species | Source |
|-------------------|------------------------|--|---|----------------------|
| Azamethiphos (OP) | 0.1 ppm | | <i>Anguilla anguilla</i> , <i>Dicentrarchus labrax</i> , <i>Oncorhynchus mykiss</i> | Intorre et al. 2004 |
| | | erratic jumping | | |
| Carbaryl (CB) | 375 and 750 µg/L | decreased swimming speed | <i>Oncorhynchus mykiss</i> | Beauvais et al. 2001 |
| | 5 µg/L (24 hr) | increased burst swimming | <i>Carassius auratus</i> | Bretau et al. 2002 |
| | 50 µg/L (24 hr) | increased sheltering and burst swimming | <i>Carassius auratus</i> | |
| Carbofuran (CB) | 500 µg/L (24 hr) | decreased swimming activity; increased sheltering, burst swimming and grouping | <i>Carassius auratus</i> | |
| | 5 µg/L (48 hr) | increased sheltering and burst swimming | <i>Carassius auratus</i> | |
| | 50 µg/L (48 hr) | increased sheltering and burst swimming | <i>Carassius auratus</i> | |
| | 500 µg/L (24 hr) | decreased swimming activity; increased sheltering, burst swimming and grouping | <i>Carassius auratus</i> | |
| | 100 µg/L | decreased attraction and buccal movements; increased sheltering | <i>Carassius auratus</i> | Saglio et al. 2003 |

| Pesticide | Dose (exposure period) | Affected Behaviour | Species | Source |
|--------------------------|---|---|-----------------------------|--------------------------------|
| Chlorpyrifos (OP) | 100 ng/mL | decreased swimming activity | <i>Danio rerio</i> | Levin et al. 2004 |
| | 0.6 µg/L | decreased swimming rate and total food stikes | <i>Oncorhynchus kisutch</i> | Sandahl et al. 2005 |
| | 1.2 µg/L | decreased swimming rate, swimming rate during feeding, and total food stikes | <i>Oncorhynchus kisutch</i> | |
| | 1.8 µg/L | decreased swimming rate, swimming rate during feeding, and total food stikes | <i>Oncorhynchus kisutch</i> | De Silva & Samayawardhena 2002 |
| | 2.5 µg/L | decreased swimming rate, swimming rate during feeding, and total food stikes; increased first food strikes | <i>Oncorhynchus kisutch</i> | |
| | 0.5 µg/L | abnormal swimming behaviour; signs of paralysis | <i>Poecilla reticulata</i> | |
| | 1 µg/L | abnormal swimming behaviour; over 50% paralysis | <i>Poecilla reticulata</i> | |
| 2 µg/L | abnormal swimming behaviour; over 50% paralysis | <i>Poecilla reticulata</i> | | |
| Chlorpyrifos-methyl (OP) | 1 mg/L | loss of equilibrium; erratic swimming; staying motionless mid-water | <i>Poecilla reticulata</i> | Selvi et al. 2005 |
| | 2.5 mg/L | staying motionless near surface then falling to aquarium bottom | <i>Poecilla reticulata</i> | |
| | 3 mg/L | loss of equilibrium; hanging verticle in water; rapid gill movement; erratic swimming; staying motionless near bottom | <i>Poecilla reticulata</i> | |

| Pesticide | Dose (exposure period) | Affected Behaviour | Species | Source |
|------------------------|------------------------|---|--|---|
| Chlorpyrifos-oxon (OP) | | tremors; flaring of gills; extension of pectoral fins; decreased escapability | <i>Gambusia affinis</i> | Carr et al. 1997 |
| | 0.08-0.16 mg/L | restlessness; erratic swimming; convulsions; loss of balance; mucous secretion | <i>Anguilla anguilla</i> <i>Lepomis macrochirus</i> | Ferrando et al. 1991 Dutta et al. 1992 |
| Diazinon (OP) | 45 µg/L | decreased optomotor behaviour; decreased following | <i>Oncorhynchus mykiss</i> | Beauvais et al. 2000 |
| | 250-1000 µg/L | decreased swimming speeds, turns, and distance, increase in tortuosity | <i>Oncorhynchus mykiss</i> | Brewer et al. 2001 |
| | 500-1000 µg/L | decreased distance travelled and swimming speed; fish swam in a more linear path | <i>Oncorhynchus tshawytscha</i> | Scholz et al. 2000 |
| | 0.1-1.0 µg/L | no change on basal foraging or swimming activity; increased activity and feeding when exposed to alarm signal | <i>Oncorhynchus tshawytscha</i> | |
| | 10.0 µg/L | no change on basal foraging or swimming activity, increased activity and feeding when exposed to alarm signal, decreased homing success | <i>Oncorhynchus tshawytscha</i> | Köprücü et al. 2006 |
| | 2 mg/L | decreased general activity and loss of equilibrium after 72 hr | <i>Silurus glanis</i> | |
| | 4-64 mg/L | decreased activity; loss of equilibrium; erratic swimming; rapid gill movement; hanging vertically in water; staying motionless on bottom | <i>Silurus glanis</i> | |
| DDVP (OP) | 1.87 mg/L | decreased food consumption | <i>Abramis brama</i> L. | Pavlov et al. 1992 |
| Folidol 600 (OP) | 2 ppm | torpor; reduction in swimming performance; remaining motionless; partial lack of reflexes | <i>Brycon cephalus</i> | de Aguiar et al. 2004 |
| | | | <i>Oncorhynchus mykiss</i> | Beauvais et al. 2000 |
| Malathion (OP) | 20-40 µg/L | decreased swimming speed and distance; increased tortuosity | <i>Oncorhynchus mykiss</i> | Beauvais et al. 2000 |
| | 20-40 µg/L | positive correlation between AChE and swimming speed, AChE and distance; negative correlation between AChE and turn rate | <i>Oncorhynchus mykiss</i> | Brewer et al. 2001 |
| | 1 and 10 µg/L | no change in rate of travel, active swimming speed or escape behaviour | <i>Sciaenops ocellatus</i> | Alvarez & Fuiman 2006 |

| Pesticide | Dose (exposure period) | Affected Behaviour | Species | Source |
|----------------------|------------------------|--|----------------------------|---------------------------------|
| Methiocarb (CB) | >5 mg/L | excessive mucus; rapid respiration rate; crowding at water surface then bottom; laterally recumbent | <i>Oncorhynchus mykiss</i> | Altinok et al. 2006 |
| Methylparathion (OP) | 4.12-5.34 mg/L | restlessness; erratic swimming; convulsions; loss of balance; mucous secretion | <i>Anguilla anguilla</i> | Ferrando et al. 1991 |
| Monocrotophos (OP) | 20.49 mg/L | dullness; loss of equilibrium; loss of feeding; erratic swimming; decreased distance traveled and swimming speed | <i>Gambusia affinis</i> | Kavitha & Venkateswara Rao 2007 |

Table A.4 Effect of temperature on the stress response in fish

| Exposure | Temperature | Effect | Species | Source |
|--|--|---|--------------------------------|------------------------|
| acclimatisation temperature | 5-35°C | fish acclimated to 5°C and 10°C had ↑ plasma cortisol levels than fish acclimated in warmer water | <i>Ictalurus punctatus</i> | Davis et al. 1984 |
| | 10-23°C | ↑ plasma glucose in fish held at 23°C | <i>Micropterus salmoides</i> | Carmichael et al. 1984 |
| Acclimatisation temperature & confinement stress | 10°C | slow ↑ in plasma cortisol and glucose | <i>Ictalurus punctatus</i> | Strange 1980 |
| | 30°C | rapid ↑ in plasma cortisol | <i>Ictalurus punctatus</i> | |
| | 20°C | moderate ↑ in plasma cortisol | <i>Ictalurus punctatus</i> | |
| | 5-30°C | fish held at 30°C had ↓ plasma glucose than fish at 5°C or 10°C | <i>Morone chrysops X</i> | Davis 2004 |
| | 5-30°C | fish held at 20°C had highest plasma cortisol levels followed by fish at 25°C, 15°C, 10°, 30°C, and 5°C | <i>Morone saxatilis</i> | |
| | 5-30°C | fish held at 30°C had highest plasma cortisol levels followed by fish at 5°C, 25°C, 21°, 16°C, and 10°C | <i>Morone chrysops X</i> | Davis & Parker 1990 |
| | 5-30°C | fish held at 5°C had highest plasma glucose levels followed by fish at 16°C, 10°C, 25°, 21°C, and 30°C | <i>Morone saxatilis</i> | |
| | 7.5, 12.5 and 21.0°C | plasma cortisol ↑ | <i>Morone saxatilis</i> | Barton & Schreck 1987 |
| | 7.5, 12.5 and 21.0°C | plasma glucose ↑, 21.0°C group had higher plasma glucose than all other groups | <i>Oncorhynchus tshawtscha</i> | |
| | 9°C | plasma cortisol ↑ | <i>Oncorhynchus tshawtscha</i> | Strange et al. 1977 |
| 20°C | plasma cortisol ↑ until 70 min where fish failed to maintain increasing cortisol production | <i>Salmo clarki clarki</i> | | |
| 5-30°C | fish held at 10°C had the smallest increase of plasma cortisol while fish held at 30°C had the highest increase of plasma cortisol | <i>Salmo clarki clarki</i> | Davis & Parker 1990 | |

| Exposure | Temperature | Effect | Species | Source |
|------------------------|--------------|--|--|-----------------------|
| decrease in water temp | 25 to 15°C | plasma glucose ↑ | <i>Chanos chanos</i> , <i>Ctenopharyngodon idella</i> | Kuo & Hsieh 2006 |
| | 25 to 10.5°C | plasma cortisol ↑ | <i>Ictalurus punctatus</i> | Weber & Bosworth 2005 |
| | 10 to 1°C | plasma cortisol ↑ - levels stayed high for at least 48 hr | <i>Oncorhynchus mykiss</i> | Barton & Peter 1982 |
| | 8 to 1°C | plasma glucose ↑ | <i>Scophthalmus maximus</i> | Staurnes 2001 |
| increase in water temp | 11 to 20°C | liver glycogen ↓ | <i>Oncorhynchus mykiss</i> | Viant et al. 2003 |
| | 0 to 10°C | plasma glucose remains steady until 9°C when it increases rapidly | <i>Pachycara brachycephalum</i> | Van Dijk et al. 1999 |
| | 20 to 31°C | plasma glucose ↓ | <i>Rhamdia quelen</i> | Lermen et al. 2004 |
| | 20 to 31°C | liver glycogen ↓ | <i>Rhamdia quelen</i> | |
| | 20 to 15°C | plasma glucose ↑ | <i>Rhamdia quelen</i> | |
| | 20 to 15°C | liver glycogen ↓ | <i>Rhamdia quelen</i> | |
| | 3 to 24°C | gradual increase in plasma glucose until 22°C then glucose levels drop | <i>Zoarcetes viviparus</i> | Van Dijk et al. 1999 |

| Exposure | Temperature | Effect | Species | Source |
|------------------------------|--------------------------|--|---|------------------------|
| low-water confinement stress | 30°C | large ↑ in plasma cortisol | <i>Morone chrysops</i> X <i>Morone saxatilis</i> | Davis 2004 |
| | 5°C | small ↑ in plasma cortisol | <i>Morone chrysops</i> X <i>Morone saxatilis</i> | |
| temperature change | change of 12°C (10-22°C) | plasma cortisol and glucose ↑ | <i>Micropterus salmoides</i> | Carmichael et al. 1984 |
| | change of 12°C (10-22°C) | plasma glucose ↑ | <i>Micropterus salmoides</i> | |
| | change of 6°C (16-22°C) | no change in plasma cortisol and glucose | <i>Micropterus salmoides</i> | |
| | change of 6°C (16-22°C) | no change in plasma glucose | <i>Micropterus salmoides</i> | |

Table A.5 Effect of temperature on AChE activity and survival during pesticide exposures

| Pesticide | Exposure | Temperature | Effect | Species | Source |
|-----------|-------------|--|--|------------------------------|----------------------|
| | | 5 - 45°C acclimation temperature of 5°C, 15°C or 25°C | Brain AChE activity increased until 35°C then brain AChE activity began to decrease | <i>Carassius auratus</i> | Hazel 1969 |
| | | acclimation temperature of 5°C, 15°C or 25°C | fish acclimated to 25°C had highest levels of brain AChE activity followed by 5°C and 15°C | <i>Carassius auratus</i> | |
| | | | no difference in AChE activity | <i>Fundulus heteroclitus</i> | |
| | | 2 - 27°C | linear ↑ in brain AChE activity | <i>Lepomis macrochirus</i> | Hogan 1970 |
| Controls | No exposure | 17.2 - 24.6°C | no change in AChE | <i>Stizostedion vitreum</i> | Phillips et al. 2002 |
| | | 5 - 20°C | linear ↑ in brain AChE activity | <i>Oncorhynchus mykiss</i> | Zinkl et al. 1987 |
| | | 20 - 25°C | slow ↑ in brain AChE activity | <i>Oncorhynchus mykiss</i> | |
| | | 25 - 35°C | plateau in brain AChE activity | <i>Oncorhynchus mykiss</i> | |
| | | >35°C | brain AChE activity ↓ | <i>Oncorhynchus mykiss</i> | |
| | | 20.1 - 30.9°C | no change in AChE activity | <i>Lepomis macrochirus</i> | Beauvais et al. 2002 |

| Pesticide | Exposure | Temperature | Effect | Species | Source |
|--------------------|------------|---------------|---|----------------------------|---------------------|
| Endosulfan (OC) | 19.78 µg/L | 13.1 - 16°C | ↓ survivors from 2.5% to 0% in small fish | <i>Oncorhynchus mykiss</i> | Capkin et al. 2006 |
| | 19.78 µg/L | 13.1 - 16°C | ↓ survivors from 42.5% to 0% in medium fish | <i>Oncorhynchus mykiss</i> | Capkin et al. 2006 |
| | 19.78 µg/L | 13.1 - 16°C | ↓ survivors from 100% to 77.5% in large fish | <i>Oncorhynchus mykiss</i> | Capkin et al. 2006 |
| Methiocarb (CB) | 5.43 mg/L | 14.9 - 17.4°C | ↑ mortality by 8.3% in large fish and 55% in small fish | <i>Oncorhynchus mykiss</i> | Altinok et al. 2006 |

Pesticide abbreviations: CB - carbamate, OC - organochlorine

| Table A.6. Species differences in AChE activity and the responses to pesticide exposure and temperature | | | | |
|---|-----------------|-------------------|---|------------------------|
| Exposure | Exposure | Dose | Species Difference | Source |
| Controls | <i>in Vitro</i> | | <i>Coregonus albula</i> had 7-fold greater brain AChE activity than <i>Perca fluviatilis</i> | Chuiko 2000 |
| | <i>in Vitro</i> | | <i>Oncorhynchus mykiss</i> had 49% less brain AChE than <i>Pimephales promelas</i> | Johnson & Wallace 1987 |
| | in water | | all species (<i>Oncorhynchus mykiss</i> , <i>Poecilia reticulata</i> , <i>Brachydanio rerio</i> and <i>Cyprinus carpio</i>) had similar levels of brain AChE activity | Kiezler et al. 1995 |
| | in water | no exposure | brain AChE activity <i>Oncorhynchus mykiss</i> > <i>E. licius</i> > <i>I. nebulosus</i> > <i>S. glanis</i> | Szabó et al. 1991 |
| | in water | no exposure | brain AChE activity <i>Pseudorasbora parva</i> > <i>C. auratus</i> , <i>O. mykiss</i> | Shaonan et al. 2004 |
| | in water | no exposure | brain AchE activity <i>Notemigonus crysoleucas</i> > <i>M. salmonides</i> , > <i>L. macrochirus</i> , > <i>G. affins</i> | Carr et al. 1997 |
| | in water | no exposure | brain AChE activity <i>Micropterus salmonides</i> > <i>G. affins</i> , > <i>L. macrochirus</i> , > <i>N. crysoleucas</i> | |
| | in water | no exposure | brain AChE activity <i>Micropterus salmonides</i> > <i>G. affins</i> , <i>N. crysoleucas</i> and <i>L. macrochirus</i> | |
| Azinphosmethyl (OP) | in water | 7.1 and 85.1 µg/L | The LC50 for <i>Sciaenops ocellatus</i> and <i>Fundulus heteroclitus</i> respectively | Van Dolah et al. 1997 |

| Exposure | Exposure | Dose | Species Difference | Source |
|---|-----------------|-----------------------------------|---|-------------------------|
| Azamethiphos | in water | 0.1 ppm | brain AChE activity <i>Anguilla anguilla</i> > <i>D. labrax</i> > <i>O. mykiss</i> | Intorre et al. 2004 |
| Chlorpyrifos (OP) | Field study | runoff from construction site | brain AChE activity <i>Micropterus salmoides</i> > <i>L. macrochirus</i> , > <i>N. crysoleucas</i> , > <i>G. affinis</i> | Carr et al. 1997 |
| Cyclohexyl methylphosphonofluoridate (OP) | <i>in Vitro</i> | 0.6-2.0 ppb | brain AChE activity <i>Carassius auratus</i> > <i>P. promelas</i> > <i>Lepomis cyanellus</i> | Weiss 1958 |
| DDVP (OP) | <i>in Vitro</i> | pI ₅₀ for each species | <i>Blicca bjoerkna</i> had 15-fold less brain AChE activity than <i>Leuciscus idus</i> | Chuiko 2000 |
| Diazinon (OP) | in water | 0.01-100 µM | brain AChE activity <i>Cyprinus carpio</i> > <i>O. mykiss</i> , > <i>P. reticulata</i> , > <i>B. rerio</i> | Kiezler et al. 1995 |
| Fenitrothion (OP) | in water | 15 to 220 µg/L | <i>Oncorhynchus mykiss</i> had greater brain AChE inhibition compared to <i>Galaxias maculatus</i> | Davies et al. 1994. |
| Malaoxon (OP) | <i>in Vitro</i> | 109-300 nM | <i>Pimephales promelas</i> was less sensitive (higher IC ₅₀) than <i>Oncorhynchus mykiss</i> | Johnson & Wallace 1987 |
| Malathion (OP) | in water | 0.25-14.5 ppm | LC ₅₀ <i>Pseudorasbora parva</i> < <i>C. auratus</i> , < <i>T. nilotica</i> , < <i>G. affinis</i> < <i>O. mykiss</i> | Shao-nan & De-fang 1996 |
| Malathion - oxidized (OP) | <i>in Vitro</i> | 0.34-0.81 µmol/l | brain AChE activity <i>Pseudorasbora parva</i> > <i>C. auratus</i> > <i>O. mykiss</i> | Shaonan et al. 2004 |
| Paraoxon (OP) | <i>in Vitro</i> | 2986-20000 nM | <i>Pimephales promelas</i> was more sensitive (lower IC ₅₀) than <i>Oncorhynchus mykiss</i> | Johnson & Wallace 1987 |
| Triazopos - oxidized (OP) | <i>in Vitro</i> | 0.042-0.16 µmol/l | brain AChE activity <i>Carassius auratus auratus</i> > <i>P. parva</i> > <i>O. mykiss</i> | Shaonan et al. 2004 |

| Exposure | Exposure | Dose | Species Difference | Source |
|-----------------------------|----------|---|---|-----------------------|
| Handling stress | in water | 30s handling stress | plasma cortisol <i>Salvelinus namaychush</i> > <i>S. fontinalis</i> , > <i>S. trutta</i> , > <i>O. mykiss</i> | Barton 2000 |
| | in water | 2hr truck transport | plasma cortisol <i>Salvelinus namaychush</i> > <i>S. fontinalis</i> > <i>O. mykiss</i> | |
| | in water | 30s handling stress | plasma glucose <i>Salmo trutta</i> > <i>S. fontinalis</i> , > <i>O. mykiss</i> , > <i>S. namaysush</i> | |
| | in water | 2hr truck transport | plasma glucose <i>Oncorhynchus mykiss</i> > <i>S. fontinalis</i> > <i>S. namaychush</i> | Jentoft et al. 2005 |
| | in water | repeatedly held out of water for 60s | Stressed <i>Perca fluviatilis</i> showed 34% lower mean body weight and <i>Oncorhynchus mykiss</i> showed 22% lower mean body weight | |
| | in water | repeatedly held out of water for 60s | <i>Perca fluviatilis</i> showed a reduced cortisol stress response whereas <i>Oncorhynchus mykiss</i> seemed to habituate to the stressor | |
| Cold shock | in water | water temperature reduced from 25 to 15°C | Plasma glucose was 2-fold higher in <i>Ctenopharyngodon idella</i> at 15°C than 25°C while plasma glucose was 5-fold higher in <i>Chanos chanos</i> at 15°C than 25°C | Kuo & Hsieh 2006 |
| Increased water temperature | in water | temperature increased from 0 to 10°C | <i>Zoarces viviparus</i> is more tolerant to heat than <i>Pachycara brachycephalum</i> | Van Dijk et al. 1999 |
| | in water | Heating 2°C/day | condition factor <i>Oncorhynchus mykiss</i> < <i>S. fontinalis</i> < <i>Salmo trutta</i> | Galbreath et al. 2004 |
| | in water | Heating 4°C/day | condition factor <i>Oncorhynchus mykiss</i> < <i>S. fontinalis</i> < <i>Salmo trutta</i> | |
| | in water | Heating 8°C/day | condition factor <i>Oncorhynchus mykiss</i> < <i>S. fontinalis</i> < <i>Salmo trutta</i> | |
| | in water | Heating 24°C/day | condition factor <i>Oncorhynchus mykiss</i> < <i>S. fontinalis</i> < <i>Salmo trutta</i> | |

| Exposure | Exposure | Dose | Species Difference | Source |
|-----------------------------|-------------|---|---|---------------|
| Decreased water temperature | in water | reduced water temperature from 8 to 1°C | Plasma glucose increased by 4-5x in <i>Scophthalmus maximus</i> L. but in <i>Hippoglossus hippoglossus</i> L. there was no change in plasma glucose | Staurnes 2001 |
| Logan River | Field study | 13-18°C | <i>Salmo trutta</i> had a higher condition factor than <i>Oncorhynchus clarkii utah</i> | McHugh 2005 |
| Logan River | Field study | 13-18°C | <i>Salmo trutta</i> reduced performance of <i>Oncorhynchus clarkii utah</i> in all temperatures but the 18°C | |