

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# exo-10,11-Dibromotricyclo[6.3.1.0<sup>2,7</sup>]-dodeca-2,4,6,9-tetraene

Kevin R. D. Johnson, Christopher O. Bender and René T. Boéré\*

Department of Chemistry and Biochemistry, University of Lethbridge, Lethbridge, AB, Canada T1K 3M4

Correspondence e-mail: boere@uleth.ca


Received 27 September 2011; accepted 10 October 2011

 Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.016;  $wR$  factor = 0.040; data-to-parameter ratio = 18.4.

The title compound,  $\text{C}_{12}\text{H}_{10}\text{Br}_2$ , is a bridged ring system based on a homobenzonorbornadiene framework. The *exo* configuration of one of the Br atoms was previously assigned *via* NMR correlations and has now been confirmed by the geometry of the solid-state structure. The compound features a Br—C—C—Br torsion angle of  $66.68(12)^\circ$ , whereby the C atoms in the calculation are respectively  $sp^3$ - and  $sp^2$ -hybridized.

## Related literature

For the structure of a closely related tribromide compound, see: Hökelek *et al.* (1991). For other similar solid-state structures based on a homobenzonorbornadiene framework, see: Daştan *et al.* (1994); Balci *et al.* (1996); Mangion *et al.* (2001). For synthesis of the title compound, see: Kitahonoki *et al.* (1969). For derivatization, see: Çakmak & Balci (1989); Bender *et al.* (2003).



## Experimental

### Crystal data

 $\text{C}_{12}\text{H}_{10}\text{Br}_2$ 
 $M_r = 314.02$ 

 Triclinic,  $P\bar{1}$   
 $a = 6.8554(5)$  Å  
 $b = 8.0926(6)$  Å  
 $c = 10.1024(7)$  Å  
 $\alpha = 78.936(1)^\circ$   
 $\beta = 78.867(1)^\circ$   
 $\gamma = 83.665(1)^\circ$ 
 $V = 538.13(7)$  Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 7.49$  mm<sup>-1</sup>  
 $T = 173$  K  
 $0.37 \times 0.36 \times 0.09$  mm

### Data collection

 Bruker APEXII CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\min} = 0.497$ ,  $T_{\max} = 0.746$   
 6108 measured reflections  
 2354 independent reflections  
 2240 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.013$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.016$   
 $wR(F^2) = 0.040$   
 $S = 1.03$   
 2354 reflections  
 128 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.58$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.45$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT-Plus (Bruker, 2008); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXTL (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

Sherry Lawson and Douglas Dolman are thanked for the sample preparation. This research was supported by the Natural Sciences and Engineering Research Council (NSERC) of Canada. The diffractometer at the University of Lethbridge X-ray Diffraction Facility was purchased with the help of NSERC and the University of Lethbridge.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2455).

## References

- Balci, M., Krawiec, M., Taskeşenligil, Y. & Watson, W. H. (1996). *J. Chem. Crystallogr.* **26**, 413–418.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bender, C. O., Dolman, D., Foesier, J. C., Lawson, S. L. & Preuss, K. E. (2003). *Can. J. Chem.* **81**, 37–44.
- Bruker (2008). APEX2, SAINT-Plus and SADABS. Bruker AXS Inc., Madison Wisconsin, USA.
- Çakmak, O. & Balci, M. (1989). *J. Org. Chem.* **54**, 181–187.
- Daştan, A., Balci, M., Hökelek, T., Ülkü, D. & Büyükgüngör, O. (1994). *Tetrahedron*, **50**, 10555–10578.
- Hökelek, T., Çakmak, O. & Balci, M. (1991). *Acta Cryst.* **C47**, 1672–1675.
- Kitahonoki, K., Takano, Y., Matsuura, A. & Kotera, K. (1969). *Tetrahedron*, **25**, 335–353.
- Mangion, D., Frizzle, M., Arnold, D. R. & Cameron, T. S. (2001). *Synthesis*, pp. 1215–1222.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2011). E67, o2975 [doi:10.1107/S1600536811041730]

**exo-10,11-Dibromotricyclo[6.3.1.0<sup>2,7</sup>]dodeca-2,4,6,9-tetraene**

**Kevin R. D. Johnson, Christopher O. Bender and René T. Boéré**

**S1. Comment**

In the title molecule (**1**) ( Fig. 1), an *exo*-configuration of Br2, which was originally assigned from NMR correlations (Kitahonoki *et al.*, 1969), has now been confirmed by the X-ray data. As such, the stereochemistry of the two enantiomers that assemble in each unit cell can be unambiguously assigned as *RRR* and *SSS* (Fig. 2).

A dihedral angle of 66.7 (1)° was calculated for the Br1—C10—C11—Br2 torsion between the *exo*-configured C11—Br2 and the *sp*<sup>2</sup> hybridized C10—Br1. Additionally, the compound contains a six-membered ring (C8—C9—C10—C11—C1—C12) that exhibits an interesting envelope-type conformation. This conformation matches that observed in other structures based on a homobenzonorbornadiene framework (Balci *et al.*, 1996). The structure reported by Hokelek *et al.* (1991) differs from **1** only by the presence of a third bromine atom attached to C12 and directed towards the cyclohexene ring (i.e. replacing H12B). However, the precision in C-C bond distances in **1** is on average five times better resulting in a considerably more reliable geometry than that reported for the tribromide.


Compound **1** has proven to be a convenient starting material for the preparation of a variety of substituted benzobarrelenes (Cakmak & Balci, 1989). For example, **1** can be readily converted to 2-bromo-3-deuteriobenzobarrelene (**2**) over several synthetic steps (Bender *et al.*, 2003) as outlined in Figure 3.

**S2. Experimental**

The dibromide (**1**) was prepared in a one-step process (Kitahonoki *et al.*, 1969) from the addition of dibromocarbene to benzonorbornadiene (**3**, Figure 3). The desired compound (**1**, m.p. 355-356 K) was obtained in 27% yield from the reaction mixture by distillation under reduced pressure and recrystallization from acetone-pentane.


**S3. Refinement**

Although hydrogen atoms were visible on the Fourier map, they were included at geometrically idealized positions with isotropic displacement parameters and refined in riding mode on their parent atoms with distance constraints: C—H = 1.00, 0.99 and 0.95 Å for methine, methylene and aromatic-type H-atoms, respectively, and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The highest residual peak had a fraction of the electron density of a single H atom and was located 0.77 Å from Br1.




**Figure 1**

A view of **1** plotted with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius.



**Figure 2**

The centrosymmetric pair of enantiomers (*SSS* and *RRR*) depicted in a packing diagram.

**Figure 3**

Synthesis of *exo*-3,4-dibromo-6,7-benzobicyclo[3.2.1]octa-2,6-diene (**1**) and its conversion to 2-bromo-3-deuteriobenzobarrelene (**2**).

### 10,11-dibromotricyclo[6.3.1.0<sup>2,7</sup>]dodeca-2,4,6,9-tetraene

#### Crystal data

C<sub>12</sub>H<sub>10</sub>Br<sub>2</sub>

*M<sub>r</sub>* = 314.02

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

*a* = 6.8554 (5) Å

*b* = 8.0926 (6) Å

*c* = 10.1024 (7) Å

$\alpha$  = 78.936 (1)°

$\beta$  = 78.867 (1)°

$\gamma$  = 83.665 (1)°

*V* = 538.13 (7) Å<sup>3</sup>

*Z* = 2

*F*(000) = 304

*D<sub>x</sub>* = 1.938 Mg m<sup>-3</sup>

Melting point = 355–356 K

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 4757 reflections

$\theta$  = 2.6–27.6°

$\mu$  = 7.49 mm<sup>-1</sup>

*T* = 173 K

Plate, colourless

0.37 × 0.36 × 0.09 mm

#### Data collection

Bruker APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 66.06 pixels mm<sup>-1</sup>

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

*T<sub>min</sub>* = 0.497, *T<sub>max</sub>* = 0.746

6108 measured reflections

2354 independent reflections

2240 reflections with *I* > 2 $\sigma$ (*I*)

*R<sub>int</sub>* = 0.013

$\theta_{\max}$  = 27.1°,  $\theta_{\min}$  = 2.1°

*h* = -8→8

*k* = -10→10

*l* = -12→12

#### Refinement

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2 $\sigma$ (*F*<sup>2</sup>)] = 0.016

*wR*(*F*<sup>2</sup>) = 0.040

*S* = 1.03

2354 reflections

128 parameters

0 restraints

Primary atom site location: heavy-atom method

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

*w* = 1/[ $\sigma^2(F_o^2) + (0.0181P)^2 + 0.3469P$ ]

where *P* = (*F<sub>o</sub>*<sup>2</sup> + 2*F<sub>c</sub>*<sup>2</sup>)/3

( $\Delta/\sigma$ )<sub>max</sub> = 0.001

$\Delta\rho_{\max}$  = 0.58 e Å<sup>-3</sup>

$\Delta\rho_{\min}$  = -0.45 e Å<sup>-3</sup>

Extinction correction: *SHELXTL* (Sheldrick,  
2008), *F<sub>c</sub>*\* = *kF<sub>c</sub>*[1 + 0.001*xF<sub>c</sub>*<sup>2</sup> $\lambda^3/\sin(2\theta)$ ]<sup>-1/4</sup>

Extinction coefficient: 0.0205 (9)

*Special details*

**Experimental.** A crystal coated in Paratone (TM) oil was mounted on the end of a thin glass fiber and cooled in the gas stream of the diffractometer Kryoflex low temperature device.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.21938 (3)	0.19786 (2)	0.215988 (17)	0.02899 (7)
Br2	0.61510 (3)	0.37752 (2)	0.314452 (18)	0.03298 (7)
C10	0.2653 (2)	0.43063 (19)	0.19221 (16)	0.0202 (3)
C9	0.2194 (2)	0.5361 (2)	0.08314 (16)	0.0210 (3)
H9	0.1728	0.4944	0.0141	0.025*
C11	0.3411 (2)	0.48460 (19)	0.30557 (16)	0.0206 (3)
H11	0.2516	0.4464	0.3944	0.025*
C7	0.0743 (2)	0.79247 (18)	0.17148 (15)	0.0193 (3)
C1	0.3474 (2)	0.67785 (19)	0.28318 (16)	0.0198 (3)
H1	0.4275	0.7135	0.3443	0.024*
C2	0.1358 (2)	0.76059 (18)	0.29894 (15)	0.0188 (3)
C12	0.4241 (2)	0.7434 (2)	0.13037 (16)	0.0224 (3)
H12A	0.4561	0.8627	0.1142	0.027*
H12B	0.5428	0.6735	0.0938	0.027*
C8	0.2415 (2)	0.72350 (19)	0.06857 (16)	0.0209 (3)
H8	0.2463	0.7860	-0.0274	0.025*
C5	-0.2360 (2)	0.9194 (2)	0.27206 (18)	0.0246 (3)
H5	-0.3634	0.9756	0.2636	0.029*
C3	0.0114 (2)	0.8056 (2)	0.41336 (16)	0.0224 (3)
H3	0.0520	0.7817	0.5002	0.027*
C4	-0.1757 (2)	0.8871 (2)	0.39833 (17)	0.0254 (3)
H4	-0.2624	0.9208	0.4755	0.030*
C6	-0.1118 (2)	0.8703 (2)	0.15684 (17)	0.0225 (3)
H6	-0.1544	0.8901	0.0707	0.027*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.04010 (11)	0.01927 (9)	0.02778 (10)	-0.00622 (7)	-0.00221 (7)	-0.00608 (6)
Br2	0.03114 (11)	0.03503 (11)	0.03414 (11)	0.01206 (7)	-0.01449 (7)	-0.00890 (7)
C10	0.0213 (7)	0.0171 (7)	0.0224 (7)	-0.0028 (6)	-0.0008 (6)	-0.0064 (6)
C9	0.0197 (7)	0.0237 (8)	0.0207 (7)	-0.0012 (6)	-0.0027 (6)	-0.0078 (6)
C11	0.0213 (7)	0.0206 (7)	0.0197 (7)	0.0022 (6)	-0.0047 (6)	-0.0040 (6)
C7	0.0222 (7)	0.0155 (7)	0.0200 (7)	-0.0026 (6)	-0.0032 (6)	-0.0028 (5)

C1	0.0188 (7)	0.0206 (7)	0.0214 (7)	-0.0012 (6)	-0.0047 (6)	-0.0057 (6)
C2	0.0197 (7)	0.0156 (7)	0.0216 (7)	-0.0022 (5)	-0.0037 (6)	-0.0036 (5)
C12	0.0204 (7)	0.0217 (8)	0.0241 (8)	-0.0029 (6)	-0.0003 (6)	-0.0041 (6)
C8	0.0233 (8)	0.0206 (7)	0.0176 (7)	-0.0005 (6)	-0.0026 (6)	-0.0021 (6)
C5	0.0189 (7)	0.0211 (8)	0.0334 (9)	-0.0001 (6)	-0.0045 (6)	-0.0048 (6)
C3	0.0254 (8)	0.0218 (8)	0.0206 (7)	-0.0027 (6)	-0.0038 (6)	-0.0052 (6)
C4	0.0239 (8)	0.0240 (8)	0.0270 (8)	-0.0019 (6)	0.0022 (6)	-0.0079 (6)
C6	0.0234 (8)	0.0199 (7)	0.0250 (8)	-0.0019 (6)	-0.0075 (6)	-0.0026 (6)

*Geometric parameters (Å, °)*

Br1—C10	1.9061 (15)	C1—H1	1.0000
Br2—C11	1.9908 (15)	C2—C3	1.382 (2)
C10—C9	1.325 (2)	C12—C8	1.539 (2)
C10—C11	1.498 (2)	C12—H12A	0.9900
C9—C8	1.517 (2)	C12—H12B	0.9900
C9—H9	0.9500	C8—H8	1.0000
C11—C1	1.541 (2)	C5—C4	1.385 (2)
C11—H11	1.0000	C5—C6	1.399 (2)
C7—C6	1.380 (2)	C5—H5	0.9500
C7—C2	1.400 (2)	C3—C4	1.397 (2)
C7—C8	1.526 (2)	C3—H3	0.9500
C1—C2	1.521 (2)	C4—H4	0.9500
C1—C12	1.541 (2)	C6—H6	0.9500
C9—C10—C11	123.77 (14)	C8—C12—C1	100.48 (12)
C9—C10—Br1	119.62 (12)	C8—C12—H12A	111.7
C11—C10—Br1	116.49 (11)	C1—C12—H12A	111.7
C10—C9—C8	119.94 (14)	C8—C12—H12B	111.7
C10—C9—H9	120.0	C1—C12—H12B	111.7
C8—C9—H9	120.0	H12A—C12—H12B	109.4
C10—C11—C1	111.60 (13)	C9—C8—C7	107.28 (12)
C10—C11—Br2	109.24 (10)	C9—C8—C12	107.79 (13)
C1—C11—Br2	108.86 (10)	C7—C8—C12	100.49 (12)
C10—C11—H11	109.0	C9—C8—H8	113.4
C1—C11—H11	109.0	C7—C8—H8	113.4
Br2—C11—H11	109.0	C12—C8—H8	113.4
C6—C7—C2	120.87 (14)	C4—C5—C6	120.84 (15)
C6—C7—C8	131.24 (14)	C4—C5—H5	119.6
C2—C7—C8	107.87 (13)	C6—C5—H5	119.6
C2—C1—C11	109.58 (12)	C2—C3—C4	118.46 (15)
C2—C1—C12	100.65 (12)	C2—C3—H3	120.8
C11—C1—C12	108.97 (12)	C4—C3—H3	120.8
C2—C1—H1	112.3	C5—C4—C3	120.70 (15)
C11—C1—H1	112.3	C5—C4—H4	119.7
C12—C1—H1	112.3	C3—C4—H4	119.7
C3—C2—C7	120.78 (14)	C7—C6—C5	118.33 (15)
C3—C2—C1	130.10 (14)	C7—C6—H6	120.8

---

C7—C2—C1	109.11 (13)	C5—C6—H6	120.8
C11—C10—C9—C8	-0.4 (2)	C2—C1—C12—C8	-42.36 (14)
Br1—C10—C9—C8	175.59 (11)	C11—C1—C12—C8	72.80 (14)
C9—C10—C11—C1	3.2 (2)	C10—C9—C8—C7	-72.51 (18)
Br1—C10—C11—C1	-172.90 (10)	C10—C9—C8—C12	34.95 (19)
C9—C10—C11—Br2	-117.19 (15)	C6—C7—C8—C9	-95.45 (19)
Br1—C10—C11—Br2	66.68 (12)	C2—C7—C8—C9	82.96 (15)
C10—C11—C1—C2	68.38 (16)	C6—C7—C8—C12	152.03 (16)
Br2—C11—C1—C2	-170.97 (10)	C2—C7—C8—C12	-29.56 (15)
C10—C11—C1—C12	-40.88 (17)	C1—C12—C8—C9	-68.19 (14)
Br2—C11—C1—C12	79.77 (13)	C1—C12—C8—C7	43.94 (14)
C6—C7—C2—C3	0.3 (2)	C7—C2—C3—C4	-1.4 (2)
C8—C7—C2—C3	-178.29 (14)	C1—C2—C3—C4	177.45 (15)
C6—C7—C2—C1	-178.77 (14)	C6—C5—C4—C3	0.4 (2)
C8—C7—C2—C1	2.62 (16)	C2—C3—C4—C5	1.1 (2)
C11—C1—C2—C3	91.68 (19)	C2—C7—C6—C5	1.1 (2)
C12—C1—C2—C3	-153.62 (16)	C8—C7—C6—C5	179.37 (15)
C11—C1—C2—C7	-89.35 (15)	C4—C5—C6—C7	-1.5 (2)
C12—C1—C2—C7	25.36 (15)		

---